

Total knee replacement may be more painful for vitamin-D deficient postmenopausal women

May 5 2021



Credit: Unsplash/CC0 Public Domain

Vitamin D is a critical part of a healthy diet. Among other benefits, it has been shown to protect against bone disease and maintain soft tissue

health. A new study suggests that it may also play a role in the degree of postoperative pain postmenopausal women experience after undergoing total knee replacement. Study results are published online today in *Menopause*.

Vitamin D deficiency is a major issue globally. It is estimated that 60% of adults have insufficient levels of the bone-building vitamin. Estrogen deficiency in perimenopausal women has been associated with decreased levels of vitamin D. A [sedentary lifestyle](#) and lack of sun exposure have also been shown to contribute to vitamin D deficiency in perimenopausal women.

In this new study, researchers sought to investigate the effect of vitamin D levels on function outcomes and risk factors of moderate to severe pain in postmenopausal women after total knee replacement. The procedure is frequently recommended for treating advanced knee osteoarthritis when nonsurgical treatment is no longer effective. Although the procedure is safe, many women experience postoperative pain.

Previous studies have sought to identify factors that play a role in determining the amount of pain women feel after undergoing knee replacement surgery. Among other factors, these studies pinpointed postmenopausal status and low estrogen levels as being associated with joint pain primarily in women aged 50 to 59 years. This new study suggests a link between vitamin D deficiency and a greater risk of postoperative pain. It identified vitamin D deficiency, smoking, and a high body mass index (BMI) as independent risk factors for moderate to severe pain after knee replacement surgery.

The new study additionally found that there was a high prevalence (67.3%) of vitamin D deficiency in postmenopausal women scheduled for [total knee replacement](#). These study results are in line with previous

studies that suggested that vitamin D deficiency is associated with the development of osteoarthritis, as well as muscle cramps, bone pain, walking difficulty, decreased bone mineral density, and fractures. The results of studies like these could provide valuable insights to clinicians evaluating postmenopausal women before major joint surgeries.

Results are published in the article "Effects of preoperative serum vitamin D levels on early clinical function outcomes and the moderate-to-[severe pain](#) prevalence in postmenopausal women after primary total knee arthroplasty."

"This study found that high body mass index, smoking, and vitamin D deficiency were independent [risk factors](#) for moderate to severe postoperative pain after knee replacement in postmenopausal women. Additionally, those with preoperative [vitamin D](#) deficiency had poorer functional outcomes. These findings highlight opportunities for clinicians to address these modifiable factors before [postmenopausal women](#) undergo joint replacement surgeries," says Dr. Stephanie Faubion, NAMS medical director.

Provided by The North American Menopause Society

Citation: Total knee replacement may be more painful for vitamin-D deficient postmenopausal women (2021, May 5) retrieved 2 May 2024 from <https://medicalxpress.com/news/2021-05-total-knee-painful-vitamin-d-deficient.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--