


Two-fifths of US seniors with COPD have poor access to pulmonary rehabilitation

May 19 2021



Seniors with Medicare have poor access to pulmonary rehab. Credit: ATS

Approximately two-fifths of Medicare beneficiaries with chronic obstructive pulmonary disease (COPD) have limited access to pulmonary rehabilitation services due to their distance from rehab centers,

according to research presented at the ATS 2021 International Conference. More than 24 million U.S. adults suffer from COPD, a smoking related disease.

Evidence has shown that pulmonary rehabilitation provides significant improvement in quality of life and functional capacity, while also reducing hospitalizations among adults with COPD. This therapeutic modality is underutilized, however. Gargya Malla, MBBS, MPH, [doctoral candidate](#), Department of Epidemiology, University of Alabama at Birmingham, and colleagues sought to quantify geographic barriers to treatment.

The researchers examined Medicare records for individuals covered between 1999 and 2018 who were diagnosed as having COPD (based on ICD9 and ICD10 codes). They identified 10,289,209 Medicare beneficiaries who had COPD. The team also established that the U.S. has 1,696 pulmonary rehabilitation centers. They then matched beneficiary zip codes with these centers.

"The 2017 National Heart, Lung, and Blood Institute (NHLBI) COPD National Action Plan emphasized the important role of pulmonary rehabilitation in COPD [patient care](#), as well as the limited access to pulmonary rehab services," noted Dr. Malla. "For the first time, we were able to quantify that limited access. The most important step to improving care for COPD patients is to increase access to pulmonary rehabilitation, a very cost-effective treatment."

Overall, the proportion of Medicare beneficiaries with COPD who, respectively, had a pulmonary rehabilitation center within 10, 15, 25 and 50 miles was 61.5 percent, 73.3 percent, 86.6 percent and 97.2 percent. A fourth of beneficiaries lived in either micropolitan, small town or [rural areas](#), which were found to have, on average, one pulmonary center per 4,300 beneficiaries with COPD. In metropolitan areas 72.8 percent of

beneficiaries with COPD had a pulmonary rehabilitation center within 10 miles, compared with 38.2 percent in micropolitan and 17.1 percent in small town and rural areas.

Previous research found that individuals who live more than 10 miles from a pulmonary rehabilitation center are less than half as likely to enroll in rehab than those who live closer.

In looking beyond the findings of her study, Dr. Malla observed that an increasing number of [pulmonary rehabilitation](#) centers are closing due to poor reimbursement from health insurers. "The COVID-19 pandemic has further exacerbated this situation," she said.

Dr. Malla suggested a number of potential solutions, such as the use of tele-rehabilitation services, expansion of supervised services such as home-based pulmonary rehab and Web-based rehab. Increased reimbursement would also help stem the closure of centers.

More information: [conference.thoracic.org/programsearch.php?sid=P6817](https://www.confERENCE.thoracic.org/programsearch.php?sid=P6817)

Provided by American Thoracic Society

Citation: Two-fifths of US seniors with COPD have poor access to pulmonary rehabilitation (2021, May 19) retrieved 24 June 2024 from <https://medicalxpress.com/news/2021-05-two-fifths-seniors-copd-poor-access.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--