


New class of compounds found to block coronavirus reproduction

June 24 2021



Transmission electron micrograph of SARS-CoV-2 virus particles, isolated from a patient. Image captured and color-enhanced at the NIAID Integrated Research Facility (IRF) in Fort Detrick, Maryland. Credit: NIAID

A human genetic mechanism hijacked by SARS-CoV-2, the coronavirus behind the COVID-19 pandemic, to help it spread also makes it vulnerable to a new class of drug candidates, a new study finds.

Led by researchers at NYU Grossman School of Medicine, a research team showed that coronavirus reproduction in infected [human cells](#) requires [chemical changes](#) made by the human protein METTL3 to RNA, a key form of genetic material. Additional human proteins involved in the recognition of modified RNA, YTHDF1 and YTHDF3, were also found to be important to the process.

Published online in *Genes and Development* on June 24, the study showed for the first time that a molecular inhibitor of METTL3, designed by Storm Therapeutics Ltd and called STM2457, dramatically reduced in cell cultures the replication of both pandemic SARS-CoV-2 and, a less severe, seasonal coronavirus, HCoV-OC43, one cause of the common cold.

"Our results represent the first time a chemical inhibitor of METTL3 has been shown to have an anti-viral effect for coronaviruses, or any virus," says senior study author Ian Mohr, Ph.D., professor in the Department of Microbiology at NYU Langone Health. "This represents a necessary step in drug development, identifies new targets, and reveals an unexpected strategy to halt the coronavirus lifecycle."

Turning virus' weaknesses against them

The current study builds on a growing understanding of gene regulation. It has long been established that sequences of As, Gs, Cs and Ts, the molecular letters in the DNA code of genes, are copied into messenger RNA (mRNA) molecules that carry the information to the machinery that determines which proteins are made. Only recently has the importance of chemical modification to mRNAs become apparent in the control of protein production. In some instances this process is controlled by the attachment of a methyl group (one carbon and three hydrogens) to an RNA chain, which turns that genetic message off.

Crucially, coronaviruses that replicate inside human cells are known to encode the complete set of their genetic instructions (their genomes) in RNA chains, raising the question of whether human RNA modification enzymes, including those that attach methyl groups, could impact the production of viral proteins that enable them to multiply.

Past work in Mohr's lab had revealed the enzymes that determine whether an A (adenosine), one of the chemical "letters" making up mRNA, is methylated at the N6 position (m6A) is important for replication of human cytomegalovirus (HCMV), with regulation by human m6A enzymes of mRNAs shaping the [immune response](#) to that virus.

In the current study, the research team showed for the first time that the reproduction of SARS-CoV-2 and HCoV-OC43 requires the action, not only of the human enzyme that installs the m6A methylation on RNA, METTL3, but human proteins that bind to this unusual arrangement of methylated RNA, YTHDF1 and YTHDF3. Remarkably, the researchers also found that the RNA genomes of both study coronaviruses contained this m6A modification.

For the next step, the NYU Langone team partnered with the UK-based Storm Therapeutics, which had run a medicinal chemistry program to develop a compound that best inhibited the action of METTL3. The current study compared the effects of the METTL3 inhibitor STM2457 and an inactive control compound, STM2120, on cultures of human lung cells infected with the seasonal coronavirus or SARS-CoV-2. The researchers then used an imaging technology to track viral infection in thousands of cells treated with different doses of STM2457.

Compared with the same concentration of the inactive control compound, the highest dose of STM2457 reduced the number of HCoV-OC43 infected cells in culture by more than 80 percent, while the same

dose of STM2457 reduced SARS-CoV-2 reproduction by more than 90 percent. Further experiments revealed that STM2457 reduced viral RNA and protein levels, but not by affecting the same human immune response mRNAs previously found to be important for HCMV.

"The inhibition of coronaviruses by this molecule is really encouraging but understanding exactly why coronaviruses need m6A RNA modification is important and might enable the design of compounds that work even better," says study first author Hannah Burgess, Ph.D., an assistant research scientist in the Department of Microbiology at NYU Langone Health.

Moving forward, the research team plans to further investigate precisely how m6A modification influences virus and host gene expression in cells infected with pandemic or seasonal coronaviruses and whether STM2457 can interfere with coronavirus replication and prevent severe disease outcomes in non-human animals.

"We went into it hoping to learn about the differences between the biology of innocuous and pandemic coronavirus infections," says co-corresponding author Angus Wilson, Ph.D., associate professor in the Department of Microbiology at NYU Langone Health. "If anything we found that both share a dependence upon the m6A methylation machinery. That creates the hope that inhibiting METTL3 may also be useful against future pandemic coronaviruses."

Provided by NYU Langone Health

Citation: New class of compounds found to block coronavirus reproduction (2021, June 24) retrieved 11 May 2024 from <https://medicalxpress.com/news/2021-06-class-compounds-block-coronavirus-reproduction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.