

# COVID-19 morbidity and mortality examined in people with rheumatic diseases

June 18 2021

---



Credit: CC0 Public Domain

The COVID-19 pandemic caused by the severe acute respiratory syndrome coronavirus 2 virus (SARS-CoV-2) is of particular concern for people with inflammatory diseases, and there are concerns that these people may be at higher risk and have poorer outcomes. However, at present the implications remain poorly understood. Population-based data from Spain show individuals with rheumatoid arthritis (RA) had an

increased risk of COVID-19 diagnosis and hospitalization compared to the general population. Similarly, data from the ARTIS database in Sweden show risks of severe COVID-19 were increased among people with inflammatory joint diseases.

Arani Vivekanantham and colleagues investigated the association between RA and the risk of COVID-19 diagnosis, hospitalization with COVID-19, and COVID-19-related [death](#). This population-based cohort study including all individuals registered in the Information System for Research in Primary Care (SIDIAP)- which covers over 80% of the population of Catalonia, Spain. This information was linked to region-wide SARS-CoV-2 testing, [hospital](#) and mortality records. Outpatient diagnoses of COVID-19, hospitalizations and deaths with COVID-19 were identified between 1st March and 6th May 2020.

A total of 5,586,565 people were identified, of which 16,344 had RA. Having RA was positively associated with being diagnosed with COVID-19, and being hospitalized with COVID-19. However, the authors did not find an association between RA and the risk of worsening from outpatient diagnosis to hospitalization or death, or from hospitalization to death.

The authors believe this is the largest study performed to date looking at COVID-19 outcomes in people with RA. Further research is needed to address factors linking RA and COVID-19 outcomes, including the presence of other comorbidities, underlying RA disease activity, and the use of immunosuppressive medications.

A second poster from Bower and colleagues looked at all-cause mortality, absolute and relative risks for severe COVID-19 in people with chronic inflammatory joint diseases, compared both over time and to the [general population](#). Using data from ARTIS—a Swedish national database -data on hospitalizations, admission to [intensive care](#), and

deaths due to COVID-19 were analyzed in 110,567 people with inflammatory joint disease, including RA, psoriatic arthritis, ankylosing spondylitis, spondyloarthritis, or juvenile idiopathic arthritis. These were compared to outcomes for 484,277 people in the general population.

In all groups, the absolute risk of death from any cause in 2020 was higher than 2015-2019, with a peak in mid-April, but the relative risks of death versus the general population remained similar.

Among people with inflammatory joint disease in 2020, the risk of hospitalization, admission to intensive care, and death due to COVID-19 was 0.5%, 0.04% and 0.1%, respectively.

Following the original abstract submission, Dr. Bower adds an update that among people with inflammatory joint disease in 2020, the risk of hospitalization, admission to intensive care, and death due to COVID-19 was 0.3%, 0.03% and 0.07%, respectively.

Provided by European Alliance of Associations for Rheumatology

Citation: COVID-19 morbidity and mortality examined in people with rheumatic diseases (2021, June 18) retrieved 27 April 2024 from <https://medicalxpress.com/news/2021-06-covid-morbidity-mortality-people-rheumatic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--