

COVID-19 weakly correlated with chilblains incidence

June 24 2021



(HealthDay)—There is a weak correlation for COVID-19 cases with

incidence of chilblains, according to a study published online June 23 in *JAMA Dermatology*.

Patrick E. McCleskey, M.D., from The Permanente Medical Group Northern California in Oakland, and colleagues conducted a [retrospective cohort study](#) to examine the potential correlation between COVID-19 incidence and chilblains. Data were included for 780 [patients](#) with chilblains reported during the pandemic.

The researchers observed a correlation for COVID-19 incidence with chilblains incidence at 207 location-months (Spearman coefficient, 0.18). Of the 456 patients with chilblains tested during the pandemic, only 3.7 percent were positive for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and only 2.0 percent were positive for SARS-CoV-2 within six weeks of chilblains diagnosis. Test results from 1.0 percent of patients were positive for immunoglobulin G antibodies to SARS-CoV-2. COVID-19, but not chilblains, disproportionately affected Latinx patients.

"Although this study found a weak geographic and temporal correlation of COVID-19 cases with chilblains cases, these findings may have resulted from [behavioral changes](#)," the authors write. "Very few cases of chilblains were positive for COVID-19."

More information: [Abstract/Full Text](#)

Copyright © 2021 [HealthDay](#). All rights reserved.

Citation: COVID-19 weakly correlated with chilblains incidence (2021, June 24) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-06-covid-weakly-chilblains-incidence.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.