


Many healthy women have labor induced for no medical reason, study shows

June 1 2021, by Hannah Dahlen, Ank de Jonge, Lilian Peters and Soo Downe



Credit: CC0 Public Domain

Pregnant women in Australia are increasingly having their labors induced rather than giving birth spontaneously—and some without good reason.

A large proportion of first time [mothers](#) (41.6%) were induced in [2018](#), when the latest national data were published, compared with 30.6% in 2010.

Our study, [published today in *BMJ Open*](#), found the induction rates tripled for [women](#) who were 37 and 39 weeks pregnant in New South Wales in the 16 years to 2016. And 15% of new mothers were induced without a medical reason listed.

Induction of labor can be life-saving in some situations. But when it's not medically indicated, it could put women and babies at unnecessary increased risk of complications.

Our induction of labor paper is out today in *BMJ Open*. Rising number of women in NSW are having induced labors without any medical reason—ABC News <https://t.co/UWgBmv9UU2>

— Hannah Dahlen AM (@hannahdahlen) [May 31, 2021](#)

Remind me, what is induction?

[Induction of labor](#) is where labor is started medically. This can be with hormones, by using a balloon-shaped catheter placed in the woman's cervix to open it up, or by breaking the bag of water around the baby.

Induction is often [recommended](#) when:

- pregnancy has gone over 41 weeks to reduce the risk of stillbirth
- the mother has [high blood pressure](#) or diabetes

- there is another significant issue threatening the health of the mother or baby.

Our research found sometimes inductions are done where there is no identified medical reason. As previous research has shown, this is especially the case in [private hospitals](#).

Sometimes women are told their baby is bigger or smaller than normal. Bigger babies may lead to more complications with the birth, and smaller babies may not be growing well. However, ultrasound can be [very inaccurate](#), and babies thought to be small or large are often a very average size at birth.

Sometimes women are sick of being pregnant and are (understandably) uncomfortable and request an induction or are offered it by doctors.

Greater intervention

Our [BMJ Open study](#) tracked almost 475,000 births in NSW between 2001 and 2016.

Of these, 69,397 (15%) had an induction of labor with no medical reason given. These women were aged 20 to 35 years, had a healthy pregnancy, and didn't smoke or have high blood pressure or diabetes.

Compared to first-time mothers who went into labor themselves, those who were induced were more likely to have:

- an instrumental birth with forceps or vacuum (28% for women who were induced vs 24% for women who gave birth spontaneously)
- a cesarean section (29% vs 14%)
- an epidural (71% vs 41%)

- an episiotomy, which is a surgical cut to the perineum, the area between the vaginal opening and the skin leading towards the anus (41% vs 30%).

In one area there were benefits for mothers who were induced: severe perineal tears were slightly lower for first time mothers (4.2% vs 4.9%) and those who had given birth previously (0.7% vs 1.2%).

Mothers having subsequent babies did not have the same high intervention rates that first time mothers did.

Another [recently published study](#) had similar findings of increased cesarean section rates for first time mothers.

Earlier inductions

We found a big rise in NSW babies being induced at what we call "[early term](#)" (37 and 38 weeks) over the 16 year period.

The number of babies born at 37 weeks' gestation tripled, while the number born at 38 weeks doubled.

Yet those last couple of weeks of being in their mother's uterus are important for the development of the child's [brain](#) and [other body systems](#) such as the lungs, and the ability to control blood sugar and body temperature.

Long-term outcomes

Previous research has suggested inducing healthy pregnant mothers after 41 weeks of pregnancy [reduces stillbirth](#) and this is what the World Health Organization recommends (we previously recommended

induction after 42 weeks).

While our research did not look at stillbirth, as all our mothers and babies were healthy when labor started, we found no difference in the rates of neonatal, infant and child death between the two groups.

Our study is one of the first to look at long-term outcomes associated with induction of labor.

Following induction of labor, [babies](#) had more trauma during birth, and were more likely to need resuscitation. This is probably partly due to being born early and/or having more surgical intervention.

Babies born after induction were more likely to be admitted to hospital with breathing difficulties and infections (ear, nose, throat, respiratory, sepsis) at a range of ages, up to 16 years.

We could only look at hospital admissions which occur when there are more serious health issues, so this does not represent visits to a GP or other community services.

Loss of control

Most women prefer to start labor spontaneously, finding induction more painful (hence more epidurals), and feeling they have less control during labor.

Women who are induced can't move around as easily due to the baby needing to be continuously monitored to pick up any signs of distress. The drugs used for induction can make the contractions very strong, reducing oxygen to the baby. Most women have intravenous fluids running, further restricting movement.

This lack of control can lead to women feeling disappointed with their birth and some may even be traumatized.

A [recent review of the evidence](#) found decisions about induction were largely made by clinicians rather than women, whose expectations and preferences were often unmet.

Time for change

There is widespread variation in [clinical practice guidelines](#) about when women should be induced.

The [World Health Organization](#) has advised against induction of labor without medical indication before 41 weeks gestation.

And this month the National Institute for Health and Care Excellence in the United Kingdom released its [draft guidance](#) for consultation, also recommending women be offered induction at 41 weeks but opening the discussion on induction up for debate.

There is no doubt induction of labor can save lives if used judiciously. But it's a major medical intervention and so should not be offered routinely before 41 weeks without discussing the risks and the potential increase in other interventions women may not anticipate.

This discussion should also include not yet knowing all the potential longer-term effects of inductions.

Most important of all, women need to be aware they can [decline or accept](#) any intervention or treatment offered or recommended by health providers and the information provided to them must be balanced, evidence based and without coercion.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Many healthy women have labor induced for no medical reason, study shows (2021, June 1) retrieved 11 May 2024 from

<https://medicalxpress.com/news/2021-06-healthy-women-labor-medical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.