

# Study links COVID-19 public health efforts to dramatic drop in COPD hospitalizations

June 14 2021

---



Credit: Pixabay/CC0 Public Domain

Researchers at the University of Maryland School of Medicine (UMSOM) analyzed data at the 13-hospital University of Maryland Medical System (UMMS) and found public health measures designed to reduce the spread of the COVID-19 virus may have fostered a substantial side benefit: Hospital admissions for chronic obstructive

pulmonary disease (COPD) were reduced by 53 percent, according to a new study published in *The American Journal of Medicine*. This is likely due to a drop in circulating seasonal respiratory viruses such as influenza.

Hospitalizations for COPD, a group of lung diseases that make it hard to breathe and get worse over time, are commonly driven by flare-ups where symptoms are triggered by such factors as tobacco smoke, air pollution and respiratory infections. Seasonal respiratory viruses, including those that cause the common cold or influenza, trigger nearly half of those flare-ups.

In the wake of a marked drop in COPD admissions during the pandemic, the researchers theorized that COVID-19 behavior changes—a mix of stay-at-home orders, social distancing, masking mandates and strict limitations on large gatherings—not only protected against COVID-19, but they may have also reduced exposure to other respiratory infections.

Conversely, they worry that the return to normal behavior may lead to more COPD flare-ups.

"Our study shows there's a silver lining to the behavior changes beyond protecting against COVID-19," said senior author Robert M. Reed, MD, UMSOM Professor of Medicine and pulmonologist at the University of Maryland Medical Center (UMMC). "If we completely eliminate masks and distancing during cold and flu season, we'll allow all those viruses that have been effectively suppressed to come raging back. There could be a lot of illness."

Prior to the COVID-19 pandemic, COPD was the fourth-leading cause of death worldwide and a leading cause of hospital admissions in the United States. The pandemic has led to significant changes in [health care delivery](#), including reduced admissions for COPD and other non-COVID

illnesses, some of which may have stemmed from patients' fear of contracting COVID in various hospital settings, as well as a shift toward telemedicine and outpatient COPD management during the pandemic.

To understand what may have occurred to reduce COPD admissions, the researchers compared weekly [hospital admissions](#) for COPD in the pre-COVID-19 years of 2018 and 2019, with admissions after the COVID-19 public health measures were instituted. At UMMS, those measures were implemented before April 1, 2020, so the investigators chose the same five-month period in each year for their comparison, April 1 to Sept. 30.

Co-lead author Jennifer Y. So, MD, UMSOM Assistant Professor of Medicine and COPD specialist at UMMC, said electronic medical records from multiple hospitals across a range of communities in the UMMS database facilitated a granular evaluation of changes over time. "We assessed a variety of possible causes that could affect COPD admissions including the presence of multiple diseases or medical conditions and the frequency of COPD exacerbations."

The database findings were correlated with data on respiratory viral trends from the U.S. Centers for Disease Control and Prevention for the period of Jan. 1, 2018, through Oct. 1, 2020.

"We found a 53 percent drop in COPD admissions throughout UMMS during COVID-19. That is substantial, but equally significant, the drop in weekly COPD admissions was 36 percent lower than the declines seen in other serious [medical conditions](#), including [congestive heart failure](#), diabetes and heart attack," said Dr. So.

As more and more people are vaccinated against COVID-19 and many of the [public health measures](#) of the past year are relaxed, the researchers warn that a full return to normal may again expose COPD

patients to the familiar seasonal triggers.

"Our study did not assess which public health components worked to tame seasonal respiratory viruses, but a simple thing like wearing a mask while riding on public transit or working from home when you're sick with a cold could go a long way to reduce virus exposure," said Dr. Reed.

Dr. So, who is from South Korea, said it is a cultural norm to wear masks during the winter in her native country. "The COVID-19 pandemic has helped a lot of people around the world become more aware of the role of masking and social distancing to reduce the spread of disease," she said.

"This is a compelling study that raises some important public health questions about protecting our most vulnerable patient populations after we are finished with the COVID-19 pandemic. I certainly think it warrants a fuller discussion," said UMSOM Dean E. Albert Reece, MD, Ph.D., MBA, University Executive Vice President for Medical Affairs and the John Z. and Akiko K. Bowers Distinguished Professor.

**More information:** Jennifer Y So et al, Decline in COPD Admissions During the COVID-19 Pandemic Associated with Lower Burden of Community Respiratory Viral Infections, *The American Journal of Medicine* (2021). [DOI: 10.1016/j.amjmed.2021.05.008](https://doi.org/10.1016/j.amjmed.2021.05.008)

Provided by University of Maryland

Citation: Study links COVID-19 public health efforts to dramatic drop in COPD hospitalizations (2021, June 14) retrieved 27 April 2024 from <https://medicalxpress.com/news/2021-06-links-covid-health-efforts-copd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.