

Study provides MIS-C treatment guidance

June 23 2021



Credit: CC0 Public Domain

An analysis conducted by a group of investigators including Tamara Bradford, MD, Associate Professor of Pediatrics at LSU Health New Orleans School of Medicine, found that children and adolescents with Multisystem inflammatory syndrome in children (MIS-C) initially treated with intravenous immune globulin (IVIG) plus glucocorticoids

had a lower risk of new or persistent cardiovascular dysfunction than IVIG alone. The research was part of the Overcoming COVID-19 Study, a nationwide collaboration of physicians at pediatric hospitals and the Centers for Disease Control and Prevention (CDC). The results were published online in the *New England Journal of Medicine*.

The researchers analyzed [surveillance data](#) on 518 children and adolescents with MIS-C who were admitted to US hospitals between March 15 and October 31, 2020. Eighty-nine (17%) received IVIG only; 241 (47%) received IVIG and glucocorticoids; 107 (21%) received IVIG, glucocorticoids, and a biologic; and 81 (16%) received other treatments, including glucocorticoids only, a biologic only, glucocorticoids and a biologic, or IVIG and a biologic. They found that [initial treatment](#) with IVIG plus glucocorticoids (103 patients) was associated with a [lower risk](#) of cardiovascular dysfunction on or after day two than IVIG alone—103 patients or 17% vs. 31%. Among those who received IVIG plus glucocorticoids, left ventricular dysfunction occurred in 8% and 17% of the patients, respectively, and shock resulting in vasopressor use in 13% and 24%. The use of adjunctive therapy was also lower among patients who received IVIG plus glucocorticoids than among those who received IVIG alone—34% vs. 70%.

"Because MIS-C cases have been sporadic, following surges of COVID-19 cases, we haven't had the benefit of randomized clinical trials of treatment strategies," notes Dr. Bradford, who practices at Children's Hospital New Orleans.

The authors write that evaluating clinical outcomes in patients with MIS-C who were treated with various immunomodulatory therapies could provide insight into their effectiveness. Until published data that define [best practices](#) are available, these data provide clinicians with additional evidence to guide treatment for MIS-C. The ongoing transmission of

SARS-CoV-2 and the emergence of variants of concern may promote continued outbreaks of MIS-C in the United States and internationally. Additional evidence-based studies are needed to examine the generalizability of these findings across a broad range of geographic regions and practice settings.

The Centers for Disease Control and Prevention defines MIS-C as a condition where different body parts can become inflamed, including the heart, lungs, kidneys, brain, skin, eyes, or gastrointestinal organs. Children with MIS-C may have a fever and various symptoms, including [abdominal pain](#), vomiting, diarrhea, neck pain, rash, bloodshot eyes, or feeling extra tired. We do not yet know what causes MIS-C. However, many children with MIS-C had the virus that causes COVID-19 or had been around someone with COVID-19.

As of June 2, 2021, a total of 4,018 patients in the US have met the MIS-C case definition, with 36 deaths. Of those, 100-149 have been in Louisiana.

More information: Mary Beth F. Son et al, Multisystem Inflammatory Syndrome in Children—Initial Therapy and Outcomes, *New England Journal of Medicine* (2021). [DOI: 10.1056/NEJMoa2102605](https://doi.org/10.1056/NEJMoa2102605)

Provided by Louisiana State University

Citation: Study provides MIS-C treatment guidance (2021, June 23) retrieved 4 May 2024 from <https://medicalxpress.com/news/2021-06-mis-c-treatment-guidance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.