

Osteoarthritis linked to higher Parkinson's disease risk

June 9 2021



Credit: Pixabay/CC0 Public Domain

A study published in *Arthritis Care & Research* has uncovered an elevated risk of Parkinson's disease in individuals with osteoarthritis.



The <u>retrospective study</u> using Taiwan's Longitudinal Health Insurance Database 2005 included 33,360 patients who were 50-64 years old and had osteoarthritis in 2002-2005. A comparison group consisted of 33,360 age- and sex- matched individuals without osteoarthritis. The osteoarthritis group had a 41% higher risk of developing Parkinson's disease, after adjustments. Patients with knee or hip osteoarthritis appeared to have a higher risk of Parkinson's disease than patients with non-knee and non-hip osteoarthritis or with uncategorized osteoarthritis.

Although the mechanisms behind a potential link between osteoarthritis and Parkinson's disease are unknown, both conditions involve inflammation.

"Coexisting osteoarthritis and Parkinson's disease can additively increase the risk of falling. Moreover, <u>osteoarthritis</u>-related mobility impairments may mask early motor symptoms of Parkinson's disease," said senior author Shin-Liang Pan, MD, Ph.D., of National Taiwan University. "Health professionals need to be alert to the potential link between these two diseases."

More information: Shih-Hao Feng et al, Osteoarthritis is associated with an increased risk of Parkinson's disease: A population-based, longitudinal follow-up study, *Arthritis Care & Research* (2021). DOI: 10.1002/acr.24708

Provided by Wiley

Citation: Osteoarthritis linked to higher Parkinson's disease risk (2021, June 9) retrieved 2 May 2024 from https://medicalxpress.com/news/2021-06-osteoarthritis-linked-higher-parkinson-disease.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.