

Secret shopper study sheds light on barriers to opioid treatment for women

June 3 2021

WOMEN FACE BARRIERS TO ACCESSING OPIOID TREATMENT, ESPECIALLY WHEN PREGNANT

Qualitative study reveals experiences of reproductive-age women accessing care

Treatment with medication improves outcomes for **women with opioid use disorder (OUD)**, yet many women - especially pregnant women - face barriers when seeking addiction treatment.

Analyzing comments from women **seeking treatment for OUD*** via phone in 10 U.S. states, qualitative research revealed:



CALL PROCESS BARRIERS

- **Calls not answered** (less than 1/2 of first attempts resulted in an appointment)
- **Calls going straight to message or automated response**
- **Calls placed on hold or transferred** (more common for pregnant callers)

"I waited almost 10 minutes for the first rep to answer the phone - she then transferred me to the scheduler which was an additional 5 minutes. The rep eventually stated that they are not accepting new patients."

"I told them I was seeking addiction treatment and four months pregnant; they put me on the classic immediate hold."

STIGMA-RELATED BARRIERS

- Disparaging comments regarding payment method, pregnancy status, addiction
- Criticism, abruptness, patronizing language and tone
- No explanation given for treatment refusal

"The receptionist very snappily said 'We don't treat pregnant women.' I barely had enough time to thank her and say goodbye before she hung up."

"Whoaaaaah, we don't do that here, I won't get into that mess again."



INSURANCE TYPE & CLINIC BARRIERS

- Cash not taken, insurance type not taken
- Requirements for previous treatment, referral, or evaluation
- Clinics not taking new patients, long wait lists




"You must first have completed an in-patient detox program [to be eligible]"

"This clinic does not accept my insurance and did not offer a self-pay option."

Not all interactions were negative - some callers were treated with kindness and respect - yet significant barriers to care were common.

There remains an urgent need to systemically improve access to medications for opioid use disorder among all women, and in particular, pregnant women.



*callers in this "secret shopper" study were trained female actors between ages 25-30 with diverse backgrounds and vocal features



VANDERBILT
Center for
Child Health Policy

@VUMCchildpolicy
www.childpolicy.org

Phillippi JC, Schulte R, Bonnet K, et al. Reproductive-Age Women's Experience of Accessing Treatment for Opioid Use Disorder: "We don't do that here." *Women's Health Issues*. Supported by NIDA R01DA045729

After a 2020 Vanderbilt University Medical Center study showed women have a difficult time accessing treatment for opioid use disorder (OUD), investigators analyzed comments received from the study's participants to further shed light on barriers to care, which included everything from long on-hold times to difficult interactions with clinic receptionists during phone calls seeking appointments. Credit: Vanderbilt University Medical Center

After a 2020 Vanderbilt University Medical Center study showed women have a difficult time accessing treatment for opioid use disorder (OUD), investigators analyzed comments received from the study's participants to further shed light on barriers to care, which included everything from long on-hold times to difficult interactions with clinic receptionists during phone calls seeking appointments.

A "secret shopper" study published in *JAMA Network Open* in 2020 used trained actors trying to get into treatment for [opioid use disorder](#) in 10 U.S. states. More than 10,000 unique "patients" were randomly assigned to be pregnant or non-pregnant and have private or Medicaid-based insurance to assess differences in the experiences of access.

The results revealed many challenges in scheduling a first-time appointment to receive medications for OUD, including finding a provider who takes insurance rather than cash. Access to treatment was even more difficult for [women](#) who are pregnant and have OUD, with pregnant women about 20% less likely to be accepted for treatment than nonpregnant women.

The trained actors provided nearly 18,000 comments describing what happened during calls, and pregnant callers and those with Medicaid

documented the most barriers in obtaining an appointment. The qualitative study is published in *Women's Health Issues*.

"Even though these trained callers didn't actually have opioid use disorder, and even though they weren't desperately trying to get help for themselves and their baby, they still felt stigmatized during these calls," said lead author Julia Phillippi, Ph.D., CNM, assistant professor and academic director of the Nurse-Midwifery program at Vanderbilt University School of Nursing. "They often felt overwhelmed by emotion because of how they were treated, both good and bad. That really says something."

Many callers experienced less than welcoming, and often stigmatizing, responses from the person answering the phone. One secret shopper reported the receptionist said, "People usually don't show up to appointments because they decide they want to have one last hoorah on dope before coming."

Callers were often placed on hold or transferred repeatedly. One caller commented, "The wait time to speak with someone is crazy," while another said, "I told them I was seeking addiction treatment and four months pregnant; they put me on the classic immediate hold."

Not all interactions were negative. One caller reported, "I would like to note that everyone I spoke with throughout the call, including the receptionist, was deeply empathetic, knowledgeable and helpful."

As the U.S. opioid crisis has grown, women have been increasingly affected. According to the National Center for Health Statistics, from 1999-2016 the rate of drug-related deaths among women jumped from 3.9 to 13.4 per thousand.

Treatment involving medications for opioid use disorder (MOUD) is

highly effective in reducing overdose death and improving quality of life. For [pregnant women](#), MOUD improve pregnancy outcomes, including [preterm birth](#).

"We're setting record levels of overdose deaths in the U.S., very likely worsened by the COVID-19 pandemic," said Stephen Patrick, MD, director of the Center for Child Health Policy at Vanderbilt University School of Medicine, associate professor of Pediatrics and Health Policy and Vanderbilt University Medical Center neonatologist. "We know medications to treat opioid use disorder reduce the risk of overdose deaths for mom and increase the likelihood that the birth is going to go to term with a higher birth weight. We should make access to treatment easy, and certainly not harder."

The authors recommend interventions to improve access to OUD care for reproductive-age women, including increasing the number of clinics in the U.S. that provide MOUD regardless of pregnancy and insurance status; more training for clinic call-takers; and addressing the issue nationally through policy reform.

More information: Julia C. Phillippi et al, Reproductive-Age Women's Experience of Accessing Treatment for Opioid Use Disorder: "We Don't Do That Here", *Women's Health Issues* (2021). [DOI: 10.1016/j.whi.2021.03.010](#)

Provided by Vanderbilt University Medical Center

Citation: Secret shopper study sheds light on barriers to opioid treatment for women (2021, June 3) retrieved 24 April 2024 from <https://medicalxpress.com/news/2021-06-secret-shopper-barriers-opioid-treatment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.