

COVID-19 anxiety syndrome still causing one in five to struggle with return to normal living

July 7 2021



Credit: CC0 Public Domain

A new survey by Kingston University in collaboration with London South Bank University (LSBU) has found that COVID-19 Anxiety Syndrome is still causing many people to struggle with reintegration into

daily life.

The new UK-wide survey of 975 people, conducted in late June, comes in the week the government set out plans to remove most COVID-19 restrictions from 19th July.

Previous research from the team involved a survey of 286 UK-based participants conducted during lockdown in February which showed a fifth of respondents were suffering from the syndrome. The latest research shows that, as restrictions have eased, a level of [anxiety](#) remains.

COVID-19 Anxiety Syndrome is a pandemic-related phenomenon that consists of forms of coping, such as attention to threat, worry, avoidance and excessive checking, that can keep people locked into a state of continuous anxiety and fear of contracting the [virus](#).

Among the key findings were that:

- One in five people scored highly on the COVID-19 Anxiety Syndrome scale.
- 40 percent strongly reported avoiding touching things in [public spaces](#) because of a fear of contracting the virus.
- 30 percent strongly reported avoiding public transport because of a fear of contracting the virus.
- 23 percent strongly reported avoiding going out to public places because of a fear of contracting the virus.
- 25 percent strongly reported paying close attention to others displaying possible symptoms of the virus.

People who had lost a family member to COVID-19 were more likely to show higher levels of COVID-19 Anxiety Syndrome. There was no particular association with age, gender, or vaccination status.

Ana Nikcevic, Professor of Psychology at Kingston University's Faculty of Business and Social Sciences, and Professor Marcantonio Spada from LSBU's Centre for Addictive Behaviours first identified the concept of COVID-19 Anxiety Syndrome in April 2020.

Their research found that some people were developing a particular set of behaviors due to their fear of the virus. They conducted the latest study in collaboration with Professor Ian Albery from LSBU's Centre for Addictive Behaviors Research.

Despite the re-opening of UK society, a significant number of people are struggling with aspects of COVID-19 Anxiety Syndrome, Professor Nikcevic said.

"Our research suggests that a significant minority continues to be avoidant and highly focused on the threat of infection, which will make return to normal daily living difficult. They will need to be supported to return to normality," she said.

The data shows that after one month of re-opening of society, many people are still struggling with aspects of the [syndrome](#), a similar figure to the previous survey conducted during full lockdown, Professor Spada added.

"This means that there are still many people who find it difficult to disengage from the COVID-19 threats which may make return to normal daily living harder as restrictions ease," he said.

"Our new findings show how vital it is that people affected by COVID-19 Anxiety Syndrome receive support. Mapping out how we will do this will become a priority for mental health service providers."

More information: Ana V. Nikčević et al, The COVID-19 anxiety

syndrome scale: Development and psychometric properties, *Psychiatry Research* (2020). [DOI: 10.1016/j.psychres.2020.113322](https://doi.org/10.1016/j.psychres.2020.113322)

Provided by Kingston University, London

Citation: COVID-19 anxiety syndrome still causing one in five to struggle with return to normal living (2021, July 7) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-07-covid-anxiety-syndrome-struggle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.