

A fermented-food diet increases microbiome diversity and lowers inflammation, study finds

July 12 2021



Credit: Unsplash/CC0 Public Domain

A diet rich in fermented foods enhances the diversity of gut microbes and decreases molecular signs of inflammation, according to researchers



at the Stanford School of Medicine.

In a clinical trial, 36 <u>healthy adults</u> were randomly assigned to a 10-week <u>diet</u> that included either fermented or <u>high-fiber foods</u>. The two diets resulted in different effects on the <u>gut microbiome</u> and the immune system.

Eating foods such as yogurt, kefir, fermented cottage cheese, kimchi and other fermented vegetables, vegetable brine drinks, and kombucha tea led to an increase in overall microbial diversity, with stronger effects from larger servings. "This is a stunning finding," said Justin Sonnenburg, Ph.D., an associate professor of microbiology and immunology. "It provides one of the first examples of how a simple change in diet can reproducibly remodel the microbiota across a cohort of healthy adults."

In addition, four types of immune cells showed less activation in the fermented-food group. The levels of 19 <u>inflammatory proteins</u> measured in <u>blood samples</u> also decreased. One of these proteins, interleukin 6, has been linked to conditions such as rheumatoid arthritis, Type 2 diabetes and chronic stress.

"Microbiota-targeted diets can change immune status, providing a promising avenue for decreasing inflammation in healthy adults," said Christopher Gardner, Ph.D., the Rehnborg Farquhar Professor and director of nutrition studies at the Stanford Prevention Research Center. "This finding was consistent across all participants in the study who were assigned to the higher fermented food group."

Microbe diversity stable in fiber-rich diet

By contrast, none of these 19 inflammatory proteins decreased in participants assigned to a high-fiber diet rich in legumes, seeds, whole



grains, nuts, vegetables and fruits. On average, the diversity of their gut microbes also remained stable. "We expected high fiber to have a more universally beneficial effect and increase microbiota diversity," said Erica Sonnenburg, Ph.D., a senior research scientist in basic life sciences, microbiology and immunology. "The data suggest that increased fiber intake alone over a short time period is insufficient to increase microbiota diversity."

The study will be published online July 12 in *Cell*. Justin and Erica Sonnenburg and Christopher Gardner are co-senior authors. The lead authors are Hannah Wastyk, a Ph.D. student in bioengineering, and former postdoctoral scholar Gabriela Fragiadakis, Ph.D., who is now an assistant professor of medicine at UC-San Francisco.

A wide body of evidence has demonstrated that diet shapes the gut microbiome, which can affect the immune system and overall health. According to Gardner, low microbiome diversity has been linked to obesity and diabetes.

"We wanted to conduct a proof-of-concept study that could test whether microbiota-targeted food could be an avenue for combatting the overwhelming rise in chronic inflammatory diseases," Gardner said.

The researchers focused on fiber and fermented foods due to previous reports of their potential health benefits. While high-fiber diets have been associated with lower rates of mortality, the consumption of fermented foods can help with weight maintenance and may decrease the risk of diabetes, cancer and cardiovascular disease.

The researchers analyzed blood and stool samples collected during a three-week pre-trial period, the 10 weeks of the diet, and a four-week period after the diet when the participants ate as they chose.



The findings paint a nuanced picture of the influence of diet on gut microbes and immune status. On one hand, those who increased their consumption of fermented foods showed similar effects on their microbiome diversity and inflammatory markers, consistent with prior research showing that short-term changes in diet can rapidly alter the gut microbiome. On the other hand, the limited change in the microbiome within the high-fiber group dovetails with the researchers' previous reports of a general resilience of the human microbiome over short time periods.

Designing a suite of dietary and microbial strategies

The results also showed that greater fiber intake led to more carbohydrates in stool samples, pointing to incomplete fiber degradation by gut microbes. These findings are consistent with other research suggesting that the microbiome of people living in the industrialized world is depleted of fiber-degrading microbes.

"It is possible that a longer intervention would have allowed for the microbiota to adequately adapt to the increase in fiber consumption," Erica Sonnenburg said. "Alternatively, the deliberate introduction of fiber-consuming microbes may be required to increase the microbiota's capacity to break down the carbohydrates."

In addition to exploring these possibilities, the researchers plan to conduct studies in mice to investigate the molecular mechanisms by which diets alter the microbiome and reduce inflammatory proteins. They also aim to test whether high-fiber and fermented foods synergize to influence the microbiome and immune system of humans. Another goal is to examine whether the consumption of fermented food decreases inflammation or improves other health markers in patients with immunological and metabolic diseases, and in pregnant women and older individuals.



"There are many more ways to target the microbiome with <u>food</u> and supplements, and we hope to continue to investigate how different diets, probiotics and prebiotics impact the <u>microbiome</u> and health in different groups," Justin Sonnenburg said.

More information: Hannah C. Wastyk et al, Gut-microbiota-targeted diets modulate human immune status, *Cell* (2021). DOI: 10.1016/j.cell.2021.06.019,

www.cell.com/cell/fulltext/S0092-8674(21)00754-6

Provided by Stanford University Medical Center

Citation: A fermented-food diet increases microbiome diversity and lowers inflammation, study finds (2021, July 12) retrieved 4 May 2024 from https://medicalxpress.com/news/2021-07-fermented-food-diet-microbiome-diversity-lowers.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.