

The rat's whiskers: Multidisciplinary research reveals how we sense texture

July 13 2021



Whisker and biomechanical measurement. (A) Rat whisker C4. The rat 's head



and location of whisker field is shown. The position of C4 is the green dot in the magnified whisker field. Conical shape, length l and distance x (cf. panel C) are shown. (Note that we repeated core measurements using also a C3 and a D3 whisker, the locations of which are indicated by grey dots) (B) Microscopic images of whisker tip (left) and base (right). (C) Experimental set up (view from the recording camera). The rotating rod is seen on top. At the bottom, the halfcylinder holding the sandpaper is shown. Experimental variables position on whisker (length) l, driving speed v, and distance x are indicated. (D) Stick-slip events in a whisker moving across a texture. Videographic analysis of whisker shape and location in x and y during one protraction (all frames of the video are shown, the sandpaper was located at distance 1 mm less than whisker length: i.e. at x=27.36 mm). Red: all frames below the driving velocity—pointing to sticking periods. Green: instances with local maximum of acceleration—pointing to slips. The first frame captured when the whisker was moving free in air (no texture) is shown for comparison (violet). Inset: Cross-correlogram of accelerations at the tip (x=24 mm, top white line) vs. that at the base (x=3 mm, bottom white line). The grey lines indicate time lag t=0 (abscissa) and correlation coefficient r=0 (ordinate). (E) Method to identify stick and slip events. Sticks (red) were found by thresholding the velocity trace (at driving speed 420° /s) and minimizing the trace below that limit. Slips (green) were found by maximizing the acceleration strips above threshold (2 standard deviations found with movement in air/no contact). On top all events are shown again aligned on the time axis. Further, a short sequence of 4 slips and 7 sticks is shown in the blow-up. Credit: *Nature* Scientific Reports

How we sense texture has long been a mystery. It is known that nerves attached to the fingertip skin are responsible for sensing different surfaces, but how they do it is not well understood. Rodents perform texture sensing through their whiskers. Like human fingertips, whiskers perform multiple tasks, sensing proximity and shape of objects, as well as surface textures.

Mathematicians from the University of Bristol's Department of



Engineering Mathematics, worked with neuroscientists from the University of Tuebingen in Germany, to understand how the motion of a whisker across a surface translates <u>texture</u> information into <u>neural signals</u> that can be perceived by the brain.

By carrying out high precision laboratory tests on a real rat whisker, combined with computation models, the researchers found that <u>whiskers</u> act like antennae, tuned to sense the tiny stick-slip motions caused by friction between the surface and the tip of the whisker.

"One of the most striking things we found both in the experiments and the theory was the thousand-fold amplification of tiny force signals perceived by the tip of the whisker to that received by the neurons at the whiskers base. Suddenly we realized that the whisker is acting like an amplifier, taking micro-scale stick-slip events and rapidly turning them into clean pulses that can be picked up and processed by the brain," said Professor Alan Champneys from the University of Bristol, co-lead of the modeling work with colleague, Dr. Robert Szalai. Dr. Thibaut Putelat carried out the detailed numerical modeling.

The research Conveyance of texture signals along a rat whisker, published in the journal *Scientific Reports* from the publisher Nature, reveals the tapering of the whisker has the effect of amplifying tiny highfrequency motions into appreciable pulse-like changes in forces and movement at the whisker follicle. In turn, the nerve cells in the follicle sense these changes and transmit them to the brain.

"It is almost as if the morphology of the whisker is designed to convey these friction-induced signals as "AC" waves on top of the "DC" motion of the whisker that conveys the information on surface proximity and hardness.

"These AC waves are too small and too rapid to be perceived by the



human eye. However, in approaching this problem in a multidisciplinary fashion, we have been able to reveal these waves with clarity for the first time," said Professor Champneys.

"The findings have implications for human touch too, where the morphology of finger-print ridges is more complex, but might similarly distinguish between AC and DC signals as our brain tries to disentangle multiple information streams about what we are feeling," said Dr. Maysam Oladazimi, who carried out the experiments as part of his Ph.D.

The findings could have far-reaching benefits including how textures could be designed to provide optimal cues for the visually impaired, for human safety operation in low light environments, or for immersive artistic installations.

"This research opens several avenues for future work. As neuroscientists, we are interested in developing a more detailed understanding of neural signaling pathways in texture discrimination, while our colleagues in Bristol are keen to explore implications for the design of future robotic sensing systems," said Professor Cornelius Schwarz, who led the experiments at the University of Tuebingen.

Professor Champneys said the research was of particular value to hapticsensing in the field of robotics, where robots literally feel their environment and is the focus of much current research, especially for robots that need to act autonomously in the dark, such as in search and rescue missions. Professor Nathan Lepora and colleagues at the Bristol Robotics Laboratory are pioneers in this field.

"This transnational interdisciplinary collaboration between experimentalists and mathematical modelers was exciting. The results from the computer models and from the laboratory experiments went hand in hand—it was only through a combination of the two that we



were able to make our breakthrough," said Professor Champneys.

More information: Maysam Oladazimi et al, Conveyance of texture signals along a rat whisker, *Scientific Reports* (2021). DOI: 10.1038/s41598-021-92770-3

Provided by University of Bristol

Citation: The rat's whiskers: Multidisciplinary research reveals how we sense texture (2021, July 13) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2021-07-rat-whiskers-</u> multidisciplinary-reveals-texture.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.