

Refining surgical treatment of long-gap esophageal atresia in children

July 26 2021, by Jessica Cerretani



Credit: Unsplash/CC0 Public Domain

The surgeons in the Esophageal and Airway Treatment Center at Boston Children's Hospital are pioneers in developing and refining innovative surgical approaches to long-gap esophageal atresia, a condition in which



a child's esophagus develops in two separate segments that can't be easily connected with surgery.

Among these techniques are jejunal interposition and the Foker process. In the former, surgeons replace the missing section of the esophagus with a section of the small bowel, specifically the jejunum. It is used primarily to treat children who have already undergone failed repair of long-gap esophageal atresia or in whom the esophagus is no longer salvageable, due to factors like caustic ingestions and esophageal perforations. It is also used in children for whom the Foker process—a technique that relies on tension-induced growth of a child's existing esophagus—is not viable.

Based on the comparatively high volume of these procedures performed at Boston Children's, our surgeons have the expertise to continually improve the management of patients with esophageal atresia. They describe these advances in two recent studies.

Supercharging in jejunal interposition

For the first paper, published in the April 2021 issue of *Surgery*, Benjamin Zendejas-Mummert, MD, MSc, and his colleagues reviewed <u>medical records</u> from 55 children who underwent jejunal interposition at Boston Children's between either 2010 and 2015 (historical cohort) or 2016 and 2019 (contemporary cohort). Specifically, they looked at the effect of "supercharging"—a type of blood flow enhancement or microvascular augmentation—on the risk of anastomotic leaks (anastomotic failure)

They found that the 41 patients in the contemporary cohort had a significantly shorter duration of medication-induced paralysis, intubation, ICU stay, and hospital stay than the 14 children in the historical cohort. They also had shorter operative times. Although the



rates of anastomotic leak were very low and similar between the two groups, no patient who had undergone a supercharged jejunal interposition experienced such a leak.

"This is a major advance, since this type of anastomosis has been historically associated with significant leak rates," explains Zendejas-Mummert. "We found that with the extra blood flow produced by supercharging, we were able to take our anastomotic leak rate to zero. Jejunal interposition remains our preferred type of esophageal replacement."

Evolution of the Foker process

A second study by Zendejas-Mummert and his colleagues looked at outcomes of the Foker process in children with long-gap EA. In the traditional Foker process, surgeons connect sutures on the two ends of the esophagus to an external traction system and then increase tension on the sutures over time. This allows the two ends to grow enough in order to be able to perform an anastomosis or staged repair. Because of the external traction system, children undergoing the Foker process must remain sedated and medically paralyzed and on a ventilator during this time. More recently, Boston Children's surgeons have developed a minimally invasive approach to the Foker process, which requires less time spent in the ICU, fewer pain medications and sedatives, and paralysis is not needed because the traction system is internal or inside the patient's chest; however not all patients are candidates for this approach.

The team compared medical records of 65 long-gap esophageal atresia patients who underwent the Foker process at Boston Children's between 2014 and 2020 (contemporary cohort) with those of 41 patients who underwent the procedure between 2005 and 2014 (historical cohort).



They found that patients in the contemporary group experienced significantly fewer leaks on traction, bone fractures, anastomotic leaks, and Foker failures than the patients in the historical cohort. Children who underwent the minimally invasive Foker process also had fewer days paralyzed, intubated, in the ICU, and in the hospital, with equivalent anastomotic outcomes. The findings were published in the February 2021 issue of the *Journal of Pediatric Surgery*.

No one-size-fits-all approach

"These results show that there is not a one-size-fits-all approach for children with long-gap esophageal atresia, but rather one must develop a custom operation for each child," says Zendejas-Mummert. "We continue to demonstrate that the best option for these children is to attempt to preserve their native esophagus. We have continued to refine and improve the outcomes of the Foker process, both in the classical open external traction and in the newer minimally invasive approach. These results further exemplify that these complex children are best cared for in centers of expertise where volume and a multidisciplinary approach to their care truly matter."

More information: Kyle Thompson et al, Evolution, lessons learned, and contemporary outcomes of esophageal replacement with jejunum for children, *Surgery* (2021). DOI: 10.1016/j.surg.2021.01.036

Wendy Jo Svetanoff et al, Contemporary outcomes of the Foker process and evolution of treatment algorithms for long-gap Esophageal Atresia, *Journal of Pediatric Surgery* (2021). DOI: <u>10.1016/j.jpedsurg.2021.02.054</u>

Provided by Children's Hospital Boston



Citation: Refining surgical treatment of long-gap esophageal atresia in children (2021, July 26) retrieved 8 May 2024 from

https://medicalxpress.com/news/2021-07-refining-surgical-treatment-long-gap-esophageal.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.