

75% of sexual assault survivors have PTSD one month later

July 20 2021

Researchers want sexual assault survivors to know that it's normal to feel awful right after the assault, but that many will feel better within three months.

In a meta-analysis published in *Trauma, Violence & Abuse*, researchers found that 81% of [sexual assault survivors](#) had significant symptoms of post-traumatic stress (PTSD) one week after the assault. One month afterward—the first point in time that PTSD can be diagnosed—75% of sexual assault survivors met criteria for the disorder. That figure dropped to 54% after three months and 41% after one year.

"One of the main takeaways is that the majority of recovery from post-[traumatic stress](#) happens in first three months," said lead author Emily Dworkin, assistant professor of psychiatry and behavioral sciences at the University of Washington School of Medicine. "We hope this will give survivors and clinicians a sense of what to expect and convey some hope."

The authors said this was the first meta-analysis of survivors' PTSD symptoms in the first year after a sexual assault. Their research underscored prior findings that PTSD is common and severe in the aftermath of sexual assault, and offered more details on the timeline for recovery.

The authors analyzed 22 studies that had assessed PTSD in sexual assault survivors over time, beginning soon after the traumatic event. The

studies cumulatively involved 2,106 sexual assault survivors.

PTSD is characterized by symptoms such as reliving a traumatic event in nightmares, [intrusive thoughts](#), or flashbacks; avoiding being reminded of the event; increases in [negative emotions](#) and decreases in [positive emotions](#); self-blame; and feeling "keyed up" or on edge, Dworkin said.

A number of proven interventions, such as Prolonged Exposure Therapy and Cognitive Processing Therapy, help people recover from sexual assault and other traumas. Dworkin said it's important for people to seek help if PTSD symptoms interfere with their functioning, no matter how much time has passed since the traumatic event.

Dworkin and Michele Bedard-Gilligan, a co-author on this study, are currently testing ways to speed the recovery process for recent survivors. One is a [smartphone app](#) that teaches evidence-based coping skills. Survivors of recent sexual assault can learn more about this and other studies by visiting thriveappstudy.com.

More information: Emily R. Dworkin et al, PTSD in the Year Following Sexual Assault: A Meta-Analysis of Prospective Studies, *Trauma, Violence, & Abuse* (2021). [DOI: 10.1177/15248380211032213](https://doi.org/10.1177/15248380211032213)

Provided by University of Washington School of Medicine

Citation: 75% of sexual assault survivors have PTSD one month later (2021, July 20) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-07-sexual-assault-survivors-ptsd-month.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.