

# Wine is safer than beer for minimizing risk of heart condition

July 29 2021, by Crispin Savage

---



Credit: CC0 Public Domain

Drinking less than six Australian-standard glasses of alcohol a week is associated with the lowest risk of developing atrial fibrillation, but not all alcohol is created equal, new research from the University of Adelaide

shows.

Atrial fibrillation (AF) is an irregular and [rapid heart rate](#) that can increase risk of stroke, [heart failure](#) and other heart-related complications. Symptoms include heart palpitations, shortness of breath and weakness.

Past research has shown that [binge drinking](#) and consumption of large amounts of alcohol increases your risk of developing atrial fibrillation but it was not clear whether consumption of low amounts of alcohol increases your risk of developing AF.

This study, published today in *JACC: Clinical Electrophysiology*, used data from the UK Biobank, a large-scale research database with [health information](#) collected from half a million UK volunteers.

Lead author Samuel Tu from the University of Adelaide's Centre for Heart Rhythm Disorders, said researchers found that those who consumed less than six Australian standard drinks of alcohol per week had the lowest risk of developing AF.

"We also found that beer and cider consumption was associated with a greater risk of atrial fibrillation, compared to red wine and white wine consumption," he said.

"Our findings suggest that responsible consumption of alcohol up to six drinks per week is safe in terms of minimizing your risk of atrial fibrillation. For those currently consuming alcohol, drinking red or white wine could potentially be a safer alternative to other types of alcoholic beverages."

"Importantly, these findings do not apply to people who already suffer from atrial fibrillation, who may find that cutting down their alcohol

consumption may reduce their symptoms."

Researchers also investigated whether the recommended number of drinks per week differed for men and women.

"A key question that we sought to answer in this study is whether the effect of alcohol is different in women and men in terms of developing [atrial fibrillation](#)," Mr Tu said.

"The answer to that is no—less than six drinks per week is the cut-off for both men and women."

**More information:** Samuel J. Tu et al, Risk Thresholds for Total and Beverage-Specific Alcohol Consumption and Incident Atrial Fibrillation, *JACC: Clinical Electrophysiology* (2021). [DOI: 10.1016/j.jacep.2021.05.013](#)

Provided by University of Adelaide

Citation: Wine is safer than beer for minimizing risk of heart condition (2021, July 29) retrieved 29 April 2024 from <https://medicalxpress.com/news/2021-07-wine-safer-beer-minimizing-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.