

Benefits of COVID-19 vaccines far outweigh very rare risk of Bell's palsy, Lancet study confirms

August 17 2021



Credit: Pixabay/CC0 Public Domain

The first large-scale population-based study on the association between the SARS-CoV-2 vaccines and Bell's palsy confirms that the beneficial

and protective effects of the vaccines far outweigh the risk of this rare adverse event. The study, published in *The Lancet Infectious Diseases* journal, finds that for every 100,000 people vaccinated with inactivated vaccine, CoronaVac, an additional 4.8 people may develop the condition.

Bell's palsy is the sudden onset of one-sided [facial paralysis](#). In the majority of cases (70%), the condition resolves itself within six months without treatment and the chance of recovery is even higher (90%) if patients receive early treatment with corticosteroids.

A small number of cases of Bell's palsy have been reported in clinical trials of mRNA COVID-19 vaccines, but so far, analyses of the association have come to conflicting conclusions. In the U.S., the Federal Drug Administration (FDA) did not consider there to be a clear causal association for the two mRNA vaccines BNT162b2 (Pfizer-BioNTech) and the mRNA-1273 (Moderna) vaccines but recommends ongoing surveillance. Acute partial facial paralysis is reported as a rare side effect of both vaccines by the European Medicines Agency. Current prescribing information for CoronaVac (an inactivated [vaccine](#)) does not list Bell's palsy as a rare adverse event, but based on the findings from the study, approval has been obtained to include information on the potential association.

Ongoing safety surveillance for any new treatment or vaccines is always recommended to understand the prevalence of rare adverse events. This is only possible with very large population studies after the initial randomised controlled trials.

In this study, researchers analysed cases of Bell's palsy related to the two approved vaccines in Hong Kong—CoronaVac and BNT162b2. The study uses data from the Hong Kong drug regulatory authority pharmacovigilance system, which includes reports of adverse events logged by health professionals throughout the territory. Cases of Bell's

palsy were included in the analysis if they occurred within 42 days of the first or second vaccine dose, within the timeframe of the study. They also conducted a nested case-control study using territory wide electronic health record database including 298 Bell's palsy cases and 1181 matched controls.

Between February 23rd, 2021 and May 4th, 2021, 28 clinically confirmed cases of Bell's palsy were identified among the 451,939 individuals who received at least a first dose of CoronaVac (equivalent to 3.61 cases per 100,000 doses administered) and 16 cases were identified among the 537,205 individuals who received at least a first dose of BNT162b2 (equivalent to 2.04 cases per 100,000 doses administered).

By analysing data from 2010-2020, the researchers estimated the background risk of Bell's palsy in Hong Kong—around 27 cases per 100,000 people, per year. Global estimates range from 15-30 cases per 100,000 people, per year. The nested case-control study found that receiving CoronaVac was associated with 2.4 times increased risk of Bell's palsy (odds ratio (OR): 2.4; 95% CI 1.4 to 4.0) whereas receiving BNT162b2 was not associated with a significantly increased risk (OR: 1.8; 95% CI: 0.9 to 3.5).

They conclude that for every 100,000 people vaccinated with CoronaVac, an additional 4.8 people may develop Bell's palsy. For BNT162b2, the increased risk was equivalent to an additional 2 cases per 100,000 people vaccinated—this finding may be attributed to underpower in the current study. Further studies with a sufficient sample size are needed to evaluate the association between Bell's palsy and BNT162b2.

"Our study suggests a small [increased risk](#) of Bell's palsy associated with CoronaVac vaccination. Nevertheless, Bell's palsy remains a rare, mostly

temporary, adverse event. All evidence to date, from multiple studies, shows that the beneficial and protective effects of the inactivated COVID-19 vaccine far outweigh any risks. Ongoing surveillance, through pharmacovigilance studies such as ours are important to calculate with increasing levels of confidence the risks of rare adverse events," says lead author Professor Ian Chi Kei Wong, The University of Hong Kong.

The authors note that they cannot conclude a causal relationship between Bell's palsy and vaccination in any individual cases from this study, and that the mechanism by which vaccination can—in very rare instances—lead to Bell's palsy remains unclear. Other studies have identified rare cases of Bell's palsy after other inactivated vaccines, such as influenza. A previous study, using the WHO Pharmacovigilance Database reported no higher risk of facial paralysis following mRNA COVID-19 vaccination than with other viral vaccines, including influenza.

The authors note that the study is limited to patients with a new diagnosis of Bell's palsy in Hong Kong, so further studies including patients with a history of Bell's palsy and patients in other regions should be done to confirm their findings. Further analyses are also needed to understand whether the risk varies by sex or age.

In a linked Comment, Professor Nicola Cirillo, University of Melbourne, Australia and Dr. Richard Doan, University of Toronto, Canada, (who were not involved in the study) write: "From a clinical, patient-oriented perspective, none of the studies published so far provide definitive evidence to inform the choice of a specific vaccine in individuals worldwide with a history of Bell's palsy. However, the data published by Wan and colleagues do offer valuable information for a rational and informed choice of COVID-19 vaccines for patients in Hong Kong, and for those in countries where both BNT162b2 and CoronaVac are

available. While waiting for conclusive evidence on vaccine-associated facial paralysis, one certainty remains: the benefit of getting vaccinated outweighs any possible risk."

More information: Eric Yuk Fai Wan et al, Bell's palsy following vaccination with mRNA (BNT162b2) and inactivated (CoronaVac) SARS-CoV-2 vaccines: a case series and nested case-control study, *The Lancet Infectious Diseases* (2021). [DOI: 10.1016/S1473-3099\(21\)00451-5](https://doi.org/10.1016/S1473-3099(21)00451-5)

Provided by Lancet

Citation: Benefits of COVID-19 vaccines far outweigh very rare risk of Bell's palsy, Lancet study confirms (2021, August 17) retrieved 12 May 2024 from <https://medicalxpress.com/news/2021-08-benefits-covid-vaccines-outweigh-rare.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.