


A new model to help clinicians predict risk of death in patients hospitalized with COVID-19

August 3 2021



Credit: Pixabay/CC0 Public Domain

Researchers from Hackensack Meridian University Medical Center and Berry Consultants, LLC, Austin, Texas have developed a new model to help clinicians predict the risk of death within 40 days in patients who are hospitalized with COVID-19 infection.

A new paper describing this retrospective, observational, multicenter cohort analysis, "Development and validation of a prognostic 40-day mortality risk model among hospitalized patients with COVID-19," was recently published in *PLOS ONE*.

The model considers six [risk factors](#): age, respiratory and oxygenation rates, and preexisting conditions such as [high blood pressure](#), coronary artery disease, or chronic kidney disease that play a role in COVID-19 deaths. Older age was determined to be the strongest predictor of death, according to health records of patients who were hospitalized with COVID-19 in a large New Jersey health network between March 1, 2020 and April 22, 2020.

"The death of every person who has passed from COVID-19 has been an enormous human tragedy, and we are working to prevent future mortality from the pandemic through vaccination, optimal care and research for new therapeutics," said Ihor S. Sawczuk, MD, FACS, regional president, Chief Research Officer, Hackensack Meridian Health northern market. "This new analysis to predict mortality in certain patients with COVID-19 was done to better understand who is most at risk for this outcome in order to do everything possible to save the lives of people who are most vulnerable to complications from this viral infection."

"It's significant that severe COVID-19 disease has occurred principally among individuals with pre-existing comorbid conditions, said Andrew Ip, MD, Division of Outcomes and Value Research, John Theurer Cancer Center at Hackensack University Medical Center, Hackensack NJ.

"Well-validated mortality models incorporating comorbidities and presenting features, similar to models used to prognosticate survival of patients in intensive care units have been lacking for COVID-19," added

Dr. Stuart Goldberg, Associate Professor of Medicine at Hackensack Meridian School of Medicine.

"Age is by far the most important risk factor in our study, with mortality risk increasing with the cube of age," said Donald A. Berry, Ph.D., a statistician with Berry Consultants. He added, "Whether this relationship is maintained for the now dominant delta variant of coronavirus SARS-CoV-2 remains to be seen."

A report from the US Centers for Disease Control and Prevention noted that 38% of COVID-19 patients had one or more underlying conditions. Patients with comorbidities are more likely to require hospitalization and intensive care support compared to individuals without additional risk factors. Moreover, high case fatality rates have been reported particularly among the elderly and people who reside in nursing homes.

The researchers developed and validated a prognostic mortality model that incorporated both pre-existing comorbidities and presenting features among people hospitalized with COVID-19 illness. Through April 22, 2020 Hackensack Meridian Healthcare's network of 13 hospitals within New Jersey had provided care to more than 3,000 COVID-19 patients and had experienced over 700 deaths. Using this cohort, the researchers presented a new model that quantifies risk of mortality within 40 days of hospitalization for COVID-19 illness.

Study Methods

The authors retrospectively reviewed the electronic health records of hospitalized patients within a 13-hospital New Jersey network between March 1, 2020 and April 22, 2020 with positive polymerase chain reaction results for SARS-CoV-2, the virus responsible for COVID-19 illness, with follow-up through May 29, 2020. With death or hospital discharge by day 40 as the primary endpoint, they used univariate

followed by stepwise multivariate proportional hazard models to develop a risk score on one-half the data set, validated on the remainder, and converted the risk score into a patient-level predictive probability of 40-day mortality.

Study Results

The study population consisted of 3123 hospitalized COVID-19 patients; median age 63 years; 60% were men; 42% had >3 coexisting conditions. 713 (23%) patients died within 40 days of hospitalization for COVID-19. From 22 [potential candidates](#), 6 factors were found to be independent predictors of mortality and were included in the risk score: age, respiratory rate ≥ 25 /minute upon hospital presentation, oxygenation

Citation: A new model to help clinicians predict risk of death in patients hospitalized with COVID-19 (2021, August 3) retrieved 24 April 2024 from <https://medicalxpress.com/news/2021-08-clinicians-death-patients-hospitalized-covid-.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--