


Dynamic stability of brain activity is altered when pondering

August 17 2021, by Zhang Nannan



Reduced stability in the default mode network and enhanced stability in the frontoparietal control network during active rumination. Credit: CHEN Xiao

People tend to keep thinking about the distress itself and the possible causes and consequences when they encounter life distress. This

psychological process is termed as "rumination" in literature. It is a common symptom and an important risk factor for major depressive disorder, but its neural mechanisms are still not clear.

Recently, a research team led by Prof. Yan Chaogan from the Institute of Psychology of the Chinese Academy of Sciences found that, the stability features of two large-scale brain networks were altered when participants were engaging active rumination.

In this study, the researchers induced participants into a continuous rumination state in a [magnetic resonance](#) imaging scanner and recorded the blood oxygen level-dependent signal during this state.

Then they characterized the temporal dynamics of the signal using a metric based on the sliding window strategy. This metric measures the overall similarity of functional connectivity architecture across all sliding windows.

After comparing the stability of rumination state and a control condition, the researchers found that the stability of the default mode network was reduced while the stability of the frontoparietal control network was enhanced during rumination.

These results indicated that the default mode network shows highly dynamic functional couplings with other [brain regions](#) to generate rumination while frontoparietal control network remain stable to exert over-constraint on the activity of default mode network.

The study, titled "Hypostability in the default mode network and hyperstability in the frontoparietal cognitive [network](#) of dynamic functional architecture during rumination," has been published in *NeuroImage* on July 24.

More information: Xiao Chen (??) et al, Hypostability in the default mode network and hyperstability in the frontoparietal control network of dynamic functional architecture during rumination, *NeuroImage* (2021). DOI: [10.1016/j.neuroimage.2021.118427](https://doi.org/10.1016/j.neuroimage.2021.118427)

Provided by Chinese Academy of Sciences

Citation: Dynamic stability of brain activity is altered when pondering (2021, August 17) retrieved 26 April 2024 from <https://medicalxpress.com/news/2021-08-dynamic-stability-brain-pondering.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.