

Epilepsy surgery may improve overall brain health

August 10 2021



Figure 1: Graphic Description of the Analysis Pipeline. (A) In addition to clinical patient data, 3D T1-weighted MRI scans were acquired for 48 patients with mesial temporal lobe epilepsy before and after temporal lobe surgery and for 37 controls. (C) Brain age was estimated for each scan using a trained



gaussian processes regression (GPR) model following tissue segmentation, vectorization, and principal components analysis (PCA)–based dimension reduction. Brain age gap estimation (BrainAGE) was computed as brain-predicted age minus actual age. BrainAGE comparisons were made between patients and control groups before and after epilepsy surgery, controlling for percentage brain volume change (SIENA; B), age, and sex (D). GM = gray matter; WM = white matter. Credit: DOI: 10.1212/WNL.00000000012289

Epilepsy surgery may be associated with improvements to overall brain health, a study led by University of Liverpool researchers has found.

The study, published in *Neurology*, the medical journal of the American Academy of Neurology, used advanced magnetic resonance imaging (MRI) techniques to inform its findings.

Corresponding author Dr. Christophe de Bezenac said: "Many clinicians remain hesitant in referring patients with refractory temporal lobe <u>epilepsy</u> (TLE) for surgery despite the known safety and efficacy. Here we provide imaging evidence to suggest that it may improve overall <u>brain</u> health."

To determine whether surgery in patients with mesial temporal lobe epilepsy (mTLE) is associated with reduced brain-predicted age as a neural marker overall brain health, the researchers compared brainpredicted and chronological age difference in patients before and after surgery to healthy controls.

Christophe de Bezenac said: "MRI analysis of brain-predicted and chronological age difference (BrainAGE) is thought to provide a surrogate marker of overall brain health. We have shown that BrainAGE is greater in patients with refractory TLE by at least seven years compared to healthy controls and that this difference is reduced after



epilepsy surgery. Our findings are in line with other work suggesting that earlier surgery may benefit patients with refractory TLE."

The researchers involved in this study are part of The Liverpool BRAIN (Brain Research Advanced Imaging Network), a clinically-oriented neuroimaging research group specializing in the application of advanced magnetic resonance imaging (MRI) techniques to understand brain impairment in neurological disorders.

More information: Christophe E. de Bézenac et al, Association of Epilepsy Surgery With Changes in Imaging-Defined Brain Age, *Neurology* (2021). DOI: 10.1212/WNL.00000000012289

Provided by University of Liverpool

Citation: Epilepsy surgery may improve overall brain health (2021, August 10) retrieved 10 May 2024 from <u>https://medicalxpress.com/news/2021-08-epilepsy-surgery-brain-health.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.