

## Low-power electrical stimulation could aid stroke rehabilitation



Synaptic remapping by electrical stimulation after stroke. (a) The connection between pre- and postsynaptic neurons are rebuilt by stroke and electrical



stimulation. After stroke newly made connections are strengthened by electrical stimulation. (b) Incoming EPSP from presynaptic neurons and additional potential provided by STES induces a postsynaptic action potential following the presynaptic action potential. (c) The upper graph, raster plot, displays the spike timing at pre- and postsynaptic neuron (Pre- and Post- respectively). The bottom graph is the simulation result of two connected neurons with STES. When STES and presynaptic activation occur asynchronously, postsynaptic neurons do not fire (left-highlighted square). When presynaptic firing and STES occur together, postsynaptic fire followed presynaptic neuron activation (right highlighted square). Credit: *Scientific Reports* 

Researchers at the Daegu Gyeongbuk Institute of Science and Technology (DGIST) in Korea have developed a new electrical brain stimulation technique that could improve the recovery of patients from strokes. Described in *Scientific Reports*, the technique uses less energy and has fewer side effects than existing methods.

A stroke is a major <u>brain</u> injury that can impair body functions, such as movement, balance and swallowing, and in order to recover, <u>brain cells</u> need to regenerate and re-establish <u>neural connections</u>. There have been various attempts to enhance this recovery process using <u>physical exercise</u> or <u>direct electrical stimulation</u> to the brain.

Electrical <u>stimulation</u>, however, can have undesirable side effects, such as speech and motor problems, caused by exciting unwanted <u>neurons</u> around the target area. In addition, <u>electrical stimulation</u> devices typically consume a lot of power, meaning they are expensive to run and require high-capacity batteries. To develop fully implantable and portable devices, these challenges must be addressed.

In response, the DGIST researchers developed a method called subthreshold electrical stimulation, through which a low-power electrical



signal can be applied to neurons at a synapse—a junction between two neurons where electric nerve impulses are transmitted.

"By reducing the power, we can minimize the impact on non-target neurons, providing a more natural and less harmful way to encourage brain recovery after a stroke," explains Kyungsoo Kim, one of the lead authors of the DGIST study.

The researchers first explored their idea using a computer simulation, before testing the technique on rats. The physical experiment involved the subthreshold electrical stimulation of post-synaptic neurons in the rats' brains. At the same time, the rats were running on a wheel which naturally stimulated the pre-synaptic neurons. This process was performed twice a day for 16 days. At the end of the treatment, they monitored the rats' behavior and tested for the presence of proteins that signify neuronal recovery.

The results showed that neither exercise nor subthreshold electrical stimulation alone were able to initiate a nerve impulse at the synapse. However, when they occurred simultaneously, the neurons reached the threshold required to fire a signal. The researchers found that the stimulated brain regions contained higher levels of specific neural proteins, indicating that neural reconstruction had taken place. They also saw improvements in the rats' behavior. Taken together, these results imply that subthreshold electrical stimulation, when combined with motor training early in rehabilitation, may strengthen connections between brain regions and aid motor recovery.

"Our approach successfully increased the survival of neural connections after a stroke, while also reducing power consumption and avoiding side effects," says Seung-Jun Yoo, another lead author of the study from DGIST.



The next step will be to test whether the technique works for strokeinduced brain damage of different severities. In the future, deep brain stimulation and other electrical treatments may be able to overcome the limitations of existing medicines to treat a wide range of different brain diseases.

**More information:** Kyungsoo Kim et al, Subthreshold electrical stimulation as a low power electrical treatment for stroke rehabilitation, *Scientific Reports* (2021). DOI: 10.1038/s41598-021-93354-x

Provided by DGIST (Daegu Gyeongbuk Institute of Science and Technology)

Citation: Low-power electrical stimulation could aid stroke rehabilitation (2021, August 10) retrieved 26 April 2024 from <u>https://medicalxpress.com/news/2021-08-low-power-electrical-aid.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.