

Non-invasive brain stimulation may reduce smoking

August 4 2021



Credit: CC0 Public Domain

A new systematic review of randomized controlled trials has found evidence that non-invasive brain stimulation may reduce smoking frequency (number of cigarettes per day) in nicotine-dependent

smokers.

Because nicotine can produce neuroadaptations in the brain that make it hard to quit smoking, researchers have sought ways to treat nicotine dependence with non-invasive brain stimulation (NIBS) techniques that counteract abnormal brain activity associated with chronic nicotine exposure.

This review, published by the scientific journal *Addiction*, looked at twelve randomized controlled trials of NIBS methods on a combined total of 710 participants with nicotine dependence. It used a network meta-analysis technique to compare the benefits and safety of various types of NIBS. Several NIBS techniques showed promising results, but the best results came from [high-frequency repetitive transcranial magnetic stimulation](#) (rTMS) of the left [dorsolateral prefrontal cortex](#) (DLPFC), which was associated with the greatest reduction in smoking frequency.

Interestingly, none of the investigated NIBS appeared to reduce the severity of either craving or [nicotine dependence](#), compared with the control groups. It may be that enhancement of DLPFC activity increases dopamine release, counterbalances the reward system, and helps patients cope with unaffected craving and withdrawal symptoms.

None of the investigated NIBS was associated with dropout rates significantly different from those of the control groups, suggesting that these kinds of treatments are well-tolerated.

Nicotine affects the release of neurotransmitters in the brain, increasing brain activity and the release of dopamine. Higher levels of dopamine bring pleasure to the smoker. Over time, chronic nicotine exposure causes the brain to produce more receptors to handle the increased [brain activity](#). When nicotine levels drop, such as when a smoker tries to quit,

activity in the brain's reward system decreases, causing withdrawal symptoms that encourage continued smoking.

More information: Tseng P-T et al, (2021) Efficacy of Noninvasive Brain Stimulation Interventions in Reducing Smoking Frequency in Patients With Nicotine Dependence: a Systematic Review and Network Meta-Analysis of Randomized Controlled Trials. *Addiction* (2021). [DOI: 10.1111/add.15624](https://doi.org/10.1111/add.15624)

Provided by Society for the Study of Addiction

Citation: Non-invasive brain stimulation may reduce smoking (2021, August 4) retrieved 20 May 2024 from <https://medicalxpress.com/news/2021-08-non-invasive-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--