

Married patients experience better outcomes following total joint arthroplasty with increased psychosocial support

August 31 2021



Credit: Unsplash/CC0 Public Domain

A recent study, "Effect of Marital Status on Outcomes Following Total Joint Arthroplasty," looked at contributing factors toward the best and

safest environment for a patient to recuperate following total knee arthroplasty (TKA) or total hip arthroplasty (THA). The study, presented at the 2021 Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS), found that, overall, married patients or those with a live-in partner experienced shorter surgical times, shorter lengths of stay (LOS) in the hospital, fewer post-operative emergency department (ED) visits, lower readmission rates and, in some cases, better patient-reported outcome measures (PROMs) than those who weren't married or lived alone.

"As [orthopaedic surgeons](#), we help restore our patients' mobility and strive to help our patients recuperate in an optimal environment," said lead researcher Ran Schwarzkopf, MD, MSc, associate chief, Division of Adult Reconstruction at NYU Langone Orthopedic Hospital. "My patient population is based in New York City and, given our urban setting, there are factors to consider such as non-elevator or walkup buildings. It's not uncommon for me to have a TKA or THA patient who lives alone on the fourth floor. Having to go up and down stairs during recovery can be a major challenge for patients—even if they don't reside in an [urban area](#)."

As director of the Adult Reconstruction Research Center at NYU Langone Orthopedic Hospital, Dr. Schwarzkopf is continually looking at factors that impact patients' recovery in order to address and mitigate obstacles, as well as optimize outcomes. One area of interest is access to physical and psychosocial (mental, emotional, social, and spiritual) support for patients following total joint arthroplasty—specifically, if a patient is married or has a live-in partner to provide a sense of safety and comfort when recuperating at home after surgery.

Dr. Schwarzkopf and his team conducted a retrospective review of patients who underwent primary THA (2,146 patients) or TKA (1,242 patients) from 2016-2020 with available PROMs and separated the

patients into two groups based on whether they were married at the time of their procedure. In the THA cohort, 1,311 (61%) patients were married and 835 (39%) were non-married. In the TKA cohort, there were 689 (55%) married patients and 553 (45%) non-married patients.

Additional factors, such as demographics (age; gender; BMI; race; smoking status; and American Society of Anesthesiologists [ASA] class, a classification of a patient's comorbidities), clinical data (surgical time; LOS; all-cause ED visits; 90-day all-cause readmissions; 90-day all-cause revisions; and discharge disposition), and PROMs were collected at various time periods.

The findings of the study include:

- Married patients had significantly shorter surgical time (88.5 vs. 93.8 minutes) and ED visits (1.5% vs. 3.8%) for THA but not for TKA.
- In both THA and TKA cohorts, married patients:
 - Had a shorter LOS (THA: 1.4 vs. 2.0 days; TKA: 2.0 vs. 2.5 days)
 - Had lower rates of 90-day all-cause readmissions (THA: 2.4% vs. 5.1%; TKA: 2.5% vs. 4.2%)
 - Were more likely to be discharged home (THA: 97.9% vs. 91.1%; TKA: 94.3% vs. 89.7%) rather than to a rehabilitation center or nursing home.
- Married patients in the THA cohort reported significantly higher PROMs scores compared to non-married patients; however, the two groups were not significantly different in the TKA cohort.

"These findings demonstrate the integral role psychosocial support can play in the recuperation period," said Dr. Schwarzkopf. "It's imperative that the multidisciplinary healthcare team conduct a proper preoperative

patient evaluation to determine any restrictions the patient faces or areas requiring additional support to help ensure the health and safety of patients post-surgery and throughout recovery."

More information: Vivek Singh et al, Effect of Marital Status on Outcomes Following Total Joint Arthroplasty, *Archives of Orthopaedic and Trauma Surgery* (2021). [DOI: 10.1007/s00402-021-03914-z](https://doi.org/10.1007/s00402-021-03914-z)

Provided by American Academy of Orthopaedic Surgeons (AAOS)

Citation: Married patients experience better outcomes following total joint arthroplasty with increased psychosocial support (2021, August 31) retrieved 11 May 2024 from <https://medicalxpress.com/news/2021-08-patients-outcomes-total-joint-arthroplasty.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--