

Why the dreaming phase matters

August 25 2021



Credit: AI-generated image (disclaimer)

Scientists have long wondered why almost all animals sleep, despite the disadvantages to survival of being unconscious. Now, researchers led by a team from the University of Tsukuba have found new evidence of brain refreshing that takes place during a specific phase of sleep: rapid eye movement (REM) sleep, which is when you tend to dream a lot.

Previous studies have measured differences in <u>blood flow</u> in the <u>brain</u>



between REM sleep, non-REM sleep, and wakefulness using various methods, with conflicting results. In their latest work, the Tsukuba-led team used a technique to directly visualize the movement of <u>red blood</u> <u>cells</u> in the brain capillaries (where nutrients and waste products are exchanged between <u>brain cells</u> and blood) of mice during awake and asleep states.

"We used a dye to make the <u>brain blood vessels</u> visible under fluorescent light, using a technique known as two-photon microscopy," says senior author of the study Professor Yu Hayashi. "In this way, we could directly observe the red blood cells in capillaries of the neocortex in nonanesthetized mice."

The researchers also measured electrical activity in the brain to identify REM sleep, non-REM sleep, and wakefulness, and looked for differences in blood flow between these phases.

"We were surprised by the results," explains Professor Hayashi. "There was a massive flow of red blood cells through the brain capillaries during REM sleep, but no difference between non-REM sleep and the awake state, showing that REM sleep is a unique state"

The research team then disrupted the mice's sleep, resulting in "rebound" REM sleep—a stronger form of REM sleep to compensate for the earlier disruption. Blood flow in the brain was further increased during rebound REM sleep, suggesting an association between blood flow and REM sleep strength. However, when the researchers repeated the same experiments in mice without adenosine A2a receptors (the receptors whose blockade makes you feel more awake after drinking coffee), there was less of an increase in blood flow during REM sleep, even during rebound REM sleep.

"These results suggest that adenosine A2a receptors may be responsible



for at least some of the changes in blood flow in the brain during REM sleep," says Professor Hayashi.

Given that reduced blood flow in the brain and decreased REM sleep are correlated with the development of Alzheimer's disease, which involves the buildup of <u>waste products</u> in the brain, it may be interesting to address whether increased blood flow in the brain capillaries during REM sleep is important for waste removal from the brain. This study lays preliminary groundwork for future investigations into the role of adenosine A2a receptors in this process, which could ultimately lead to the development of new treatments for conditions such as Alzheimer's disease.

The research was published in Cell Reports.

More information: Chia-Jung Tsai et al, Cerebral capillary blood flow upsurge during REM sleep is mediated by A2a receptors, *Cell Reports* (2021). DOI: 10.1016/j.celrep.2021.109558

Provided by University of Tsukuba

Citation: Why the dreaming phase matters (2021, August 25) retrieved 26 April 2024 from <u>https://medicalxpress.com/news/2021-08-phase.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.