

Study shows missed potential for healthy life after a heart attack

August 26 2021



Credit: CC0 Public Domain

Adherence to lifestyle advice and medications could add seven healthy years of life after a heart attack, according to a study presented at ESC Congress 2021.

"Most [heart attack patients](#) remain at high risk of a second attack one year later," said study author Dr. Tinka Van Trier of Amsterdam University Medical Centre, the Netherlands. "Our study suggests that improving both lifestyles and medication use could lower this risk, with a gain in many years of life without a cardiovascular event."

The INTERHEART study previously demonstrated that 80–90% of the risk of a [heart](#) attack can be modified by managing factors such as smoking, unhealthy diet, abdominal obesity, inadequate physical activity, hypertension, diabetes and raised [blood lipid levels](#). Such management consists of two main strategies: lifestyle change and medication.

However, the RESPONSE studies showed that adequate levels of these risk factors are seldom reached after a heart attack, even in programmes aiming to help patients improve their lifestyles and optimise their medication. Therefore, "residual risk", i.e. the risk for another heart attack that is left after conventional treatment, is high to very high in a large number of patients. Dr. Van Trier said: "This study was conducted to quantify this residual risk and estimate the extent to which it could be lowered with optimal management."

The study pooled data from 3,230 patients that had a heart attack or received a stent or bypass surgery. The average age was 61 years and 24% were women. At an average of one year after the cardiac event, nearly one in three (30%) continued smoking, 79% were overweight, and 45% reported insufficient physical activity. Just 2% reached treatment targets for blood pressure, LDL ("bad") cholesterol, and glucose levels—with 40% having [high blood pressure](#) and 65% having high LDL cholesterol. However, use of preventive medications was common: 87% used antithrombotic medications, 85% took lipid lowering drugs and 86% were on blood pressure lowering drugs.

Using the SMART-REACH model, the researchers calculated the

lifetime risk of a heart attack, stroke, or death from cardiovascular disease and estimated changes in healthy years, i.e. cardiac event-free, when lifestyle or medication was changed or optimised. The model incorporates the following treatment targets: 1) not smoking; 2) antithrombotic therapy with two antiplatelet drugs; 3) lipid lowering medication (high-intensity statin, ezetimibe and PCSK9 inhibitor); 4) systolic [blood pressure](#) below 120 mmHg; 5) if diabetic, use of GLP1-agonist and SGLT2 inhibitor and controlled blood sugar (HbA1c less than 48 mmol/mol).

Dr. Van Trier explained: "The model does not incorporate all [lifestyle advice](#) since quantitative data are lacking to calculate gains in healthy life years. But that does not mean that recommendations to eat healthily, maintain a normal weight, and do regular physical activity are less important to reduce your risk."

The estimated average residual lifetime risk was 54% – meaning that half would have a heart attack, stroke, or die from cardiovascular disease at some point during their life. If the treatment of patients in the study was optimised to meet all targets in the model, the average risk would drop to 21% (one in five patients).

Dr. Van Trier said: "The findings show that despite current efforts to reduce the likelihood of new events after a [heart attack](#), there is considerable room for improvement. Our analysis suggests that the risk of another [cardiovascular event](#) could, on average, be halved if therapies were applied or intensified. For individual patients, this would translate into gaining an average of 7.5 event-free years."

More information: Abstract title: Modifiable lifetime risk for recurrent major cardiovascular events: observations in a contemporary pooled cohort.

Provided by European Society of Cardiology

Citation: Study shows missed potential for healthy life after a heart attack (2021, August 26)
retrieved 5 May 2024 from

<https://medicalxpress.com/news/2021-08-potential-healthy-life-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--