

Tick bites: Every year is a bad tick year

August 19 2021, by Jory Brinkerhoff



Black-legged ticks carry Lyme disease, which continues to spread widely across the United States. Credit: CDC/Michael Levin

It's summer, a time to hike, garden, vacation—and to be on the lookout for ticks.


From Lyme disease to lesser-known illnesses like <u>Heartland virus disease</u>, <u>ehrlichiosis</u> and <u>Colorado tick fever</u>, tick-borne disease cases are <u>increasing rapidly</u> in the United States.

In 2017, <u>59,349 cases</u> were reported to the U.S. Centers for Disease



Control and Prevention, an all-time high. Yet, this represents just a fraction of infections because those who don't exhibit symptoms or fail to seek treatment remain uncounted. A recent report estimated nearly a half-million Lyme disease cases per year in the U.S., with numbers more than doubling from 2004 to 2016.

As a biologist who <u>studies tick-borne disease</u>, I am asked each spring and summer whether it will be a bad year for ticks. The answer: It is never a good year for ticks. There may be relatively few of certain <u>species</u> and many of other types. Different species of ticks live in different environments. Many factors influence numbers, from <u>dwindling</u> <u>biodiversity</u> and <u>ecological change</u> to the <u>changing climate</u>. But every year, the time to be most vigilant is early spring through late fall.



Michigan, Wisconsin and the Northeast are hot spots for tick-borne disease in the U.S. Credit: CDC

Different species, different patterns




There are at least <u>seven tick species</u> in North America that commonly bite and infect humans and animals with numerous <u>diseases</u>. But there are others, too. Over the past two decades, seven new tick-borne germs have been identified in the U.S., including a newly discovered Lyme disease <u>bacterium</u> found in the Upper Midwest and Bourbon virus, discovered in Bourbon County, Kansas.

Some regions, like where I work in Richmond, Virginia, are home to multiple human-biting species, each with its own suite of pathogens and habitat preferences. Black-legged ticks, which spread Lyme and other diseases, are of greatest concern. They are common in forests across the Eastern U.S.. The bite of an infected American dog tick, which prefer grassy areas, can infect people and dogs with Rocky Mountain spotted fever. Aggressive Lone Star ticks, which can transmit ehrlichiosis and tularemia, thrive in many habitats across the eastern U.S., and can survive hot, dry conditions.

Complex life cycles

These parasitic arthropods are more closely related to mites, spiders and scorpions than to insects. Ticks spend <u>most of their time</u> on the ground in leaf litter or vegetation, undergoing a <u>four-stage metamorphosis</u>.





The life cycle of a black-legged tick is generally two years. Credit: CDC

Eggs hatch into six-legged larvae and attach to a host. Then they drop off and molt into eight-legged nymphs, find a host and grow into reproductive adults. Much of their lives are spent waiting, inactive, for warmer or more humid weather to continue development, or on the hunt for their next meal.

Mortality is high. If just 10% of ticks survived each life stage, it would take 2,000 eggs to produce a pair of reproductive adults. Small changes



in survival can affect populations for years.

Most of the hard-bodied, blood-feeding ticks that carry disease take only three bloodmeals during their entire <u>two- to three-year life cycle</u>—meals that allow them to molt into the next stage, or to lay eggs.

They <u>feed</u> on mammals, birds, reptiles and amphibians. Most ticks prefer a different host at each stage of their life, finding it by detecting an animal's breath or smell, sensing body heat, moisture or vibrations.

The numbers of <u>available hosts</u> may be a key factor in tick abundance, which is sometimes influenced by natural cycles. For example, during <u>"mast" years</u> when acorns are plentiful, white-footed mice populations grow along with black-legged ticks that feed on them, and <u>Lyme disease cases</u> also tend to rise.

Expanding territory

Predicting tick numbers grows harder as many species expand their ranges. Altered ecosystems play a substantial role. Lyme disease became epidemic when mice that carry the bacterium proliferated and deer were reintroduced for hunting in the 1900s after a steep decline; deer act as hosts for adult black-legged ticks that spread the disease. Migrating birds have also helped <u>disperse ticks</u> along the Atlantic flyway.

<u>Changing climate</u>, with shorter, milder winters, may increase tick survival, creating larger populations. Shorter, warmer winters have allowed some species to move northward. The two Lyme-spreading tick species now live in at least <u>43 states</u>. <u>Gulf Coast</u> ticks have spread north to Delaware and Illinois, and the Lone Star tick <u>may soon reach Canada</u>.

However, warming trends may have mixed impacts. Ticks need moisture as well as blood to survive; hot, dry weather kills some species, but not



others.

The forecast

While researchers have identified why their numbers change over space and time, predicting risk is difficult. But we do know that tick-borne diseases will continue to be a human and veterinary health threat. Ancient ticks once <u>fed on dinosaurs</u>. Scientists discovered fossilized ticks, some 15 million years old, that carried the Borrelia bacteria that causes Lyme <u>disease</u>, showing that it <u>existed long before humans</u>.

From a public health perspective, the most important question about ticks is not whether a given year will be particularly bad in terms of <u>tick</u> numbers, but what can be done to <u>reduce the risk</u> of encountering these parasites to avoid illness.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Tick bites: Every year is a bad tick year (2021, August 19) retrieved 18 April 2024 from https://medicalxpress.com/news/2021-08-year-bad.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.