

Deficits may remain after 7-day recovery from 10 days of insufficient sleep

September 1 2021



Credit: CC0 Public Domain

After 7 days of recovery from a 10-day period of deficient sleep, participants in a small study had recovered their pre-sleep deprivation reaction speed, but had not fully recovered on any other measures of



function. Jeremi Ochab of the Jagiellonian University in Kraków, Poland, and colleagues present these findings in the open-access journal *PLOS ONE* on September 1, 2021.

Sleep deficiency is well known to negatively impact human functioning. For example, it is associated with deficits in attention and memory, as well as increased risk of car accidents, heart problems, and other medical issues. However, while some research has addressed <u>recovery</u> after <u>chronic sleep deprivation</u>, it has been unclear how much time is needed to fully recover from prolonged periods of deficient sleep.

To shed more light on this topic, Ochab and colleagues conducted a small study with several <u>healthy adults</u> who underwent 10 days of purposeful sleep restriction followed by 7 recovery days of unrestricted sleep. Participants completed the study in their normal day-to-day environments and wore wrist sensors to monitor daily patterns of sleep and activity. They also underwent daily electroencephalography (EEG) to monitor <u>brain activity</u>, and they answered daily questions (Stroop tasks) to measure reaction times and accuracy.

After 7 days of recovery, the participants had not yet returned to presleep deprivation performance on most measures of functioning. These included several EEG measures of brain activity, rest-versus-activity patterns captured by wrist sensors, and accuracy on Stroop tasks. Only their reaction times had recovered to baseline levels.

While the researchers note that it is difficult to compare these results with other studies that employed different methods, the findings contribute new insights into recovery from chronic sleep loss. Future research could expand to a greater number of participants, investigate longer recovery periods, and disentangle the order in which different functions return to normal.



The authors add: "The investigation of the recovery process following an extended period of sleep restriction reveal that the differences in behavioral, motor, and neurophysiological responses to both sleep loss and recovery."

More information: Ochab JK, Szwed J, Oleś K, Bereś A, Chialvo DR, Domagalik A, et al. (2021) Observing changes in human functioning during induced sleep deficiency and recovery periods. *PLOS ONE* 16(9): e0255771. doi.org/10.1371/journal.pone.0255771

Provided by Public Library of Science

Citation: Deficits may remain after 7-day recovery from 10 days of insufficient sleep (2021, September 1) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2021-09-deficits-day-recovery-days-insufficient.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.