

Doctors often miss signs of type 1 diabetes in kids

September 21 2021



(HealthDay)— Potentially dangerous symptoms of type 1 diabetes in

children are not always immediately recognized by primary care providers, new research suggests.

In type 1 diabetes, the pancreas fails to make enough insulin, the hormone that helps regulate blood sugar used for energy by cells. Between 5% and 10% of cases of diabetes are type 1, which often first surfaces in childhood.

The Swedish study, published in the September issue of the journal [*Pediatric Diabetes*](#), suggests that key signs of type 1 diabetes—thirst, excessive urination, fatigue, weight loss—are often not diagnosed as type 1 disease in children brought to primary care physicians.

The research was led by Johan Wersäll, of the University of Gothenburg. His team surveyed the caregivers of 237 children and teens under the age of 19. All had been admitted to the hospital with new-onset type 1 diabetes and diabetic ketoacidosis between 2015 and 2017.

Diabetic ketoacidosis (DKA) is a potentially dangerous condition that can occur when diabetes goes untreated. It occurs when, starved of [blood sugar](#), cells turn to burning fat to produce energy instead. A side effect of this is the dangerous buildup of chemicals in the blood called ketones—left unchecked, this can trigger coma and even death.

"The only way to avoid DKA in patients with new-onset type 1 diabetes is to initiate insulin therapy urgently," the researchers noted.

In the study, the investigators found that 39% of parents had had suspicions of new-onset diabetes before they brought their child to [medical care](#).

Wersäll's group looked at 112 cases where parents had first brought a child they thought was ill to a primary care physician, and then ended up

bringing the same child to a hospital emergency ward within the next four weeks.

In 64 of the cases, the [primary care physicians](#) did spot type 1 symptoms and quickly referred the child to hospital emergency care, the Swedish team reported. But in 48 other cases these referrals were delayed.

"Classic symptoms of thirst, polyuria [excessive urination], tiredness and [weight loss](#) were the most common in both groups," they noted. In many cases, despite these classic symptoms, "urinary glucose [sugar] or blood glucose levels were not tested," the team noted.

In cases with a delayed diagnosis, "blood glucose or urinary glucose had been tested at the primary health care center" only 42% of the time, despite kids showing classic type 1 symptoms, the study found.

And in many cases where elevated [blood glucose levels](#) were confirmed, children were still "not directly referred to hospital," the researchers found.

The bottom line, according to the investigators: "as many as 43% of these children were not immediately referred to a pediatric emergency ward, indicating a substantial doctor's delay. Improved knowledge and general awareness of [diabetes](#) symptoms among both caregivers and among medical professionals working in the primary health care sector are paramount in improving this situation and preventing [diabetic ketoacidosis](#)."

More information: Find out more about type 1 diabetes at the [American Diabetes Association](#).

Copyright © 2021 [HealthDay](#). All rights reserved.

Citation: Doctors often miss signs of type 1 diabetes in kids (2021, September 21) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-09-doctors-diabetes-kids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.