


# Research proves feasibility of new cancer treatment approach

September 1 2021

---



Credit: CC0 Public Domain

Researchers from Cleveland Clinic and OhioHealth have published results from a first-in-human trial evaluating feasibility of Temporally Feathered Radiation Therapy (TFRT), which is designed to reduce the toxicity of radiation treatment.

The research, appearing in *Radiotherapy & Oncology*, proved TFRT's feasibility in clinical workflow. Additionally, assessments of treatment toxicities and radiation dosage comparisons to a standard radiotherapy plan were described.

TFRT is a novel technique for the planning and optimization of radiotherapy that considers the nonlinear aspects of normal tissue repair to manage toxicity. The technique is designed to reduce radiation-induced toxicities by optimizing the time through which radiation is delivered and consequently improves normal tissue recovery. Radiation-induced toxicity is a major contributor to impacting a patients' quality of life and often a dose-limiting factor in the treatment of cancer with radiation therapy.

"We are excited that the complex TFRT technique, which uses current planning systems, was able to be delivered safely and in a standard clinical workflow," said Jacob G. Scott, M.D., DPhil, a Cleveland Clinic radiation oncologist and the inventor of TFRT. "A larger trial with toxicity as the primary endpoint will allow us to truly study the efficacy of the approach. Collaboration will be important as we work to integrate TFRT into planning systems to expand automation and wider adoption into clinical practice."

In this study, five patients with head and neck squamous cell carcinoma

were treated with TFRT. The primary endpoint was feasibility of TFRT planning as defined by radiation start within 15 days of treatment planning. Secondary endpoints included estimates of toxicity.

The primary endpoint was met when patients were successfully treated with TFRT techniques without causing delays in radiation commencement. For patients who received TFRT, the median time from treatment planning to radiation start was 10 business days—not outside standard timelines. The average time required for radiation planning was six days. The organs feathered included oral cavity, each submandibular gland, each parotid gland, supraglottis, and posterior pharyngeal wall. There were no significant deviations from standard planning, and toxicity was no more than expected—but we expect a larger trial to show reductions in toxicity without effecting cure rates.

"Radiation oncologists and physicists have made amazing advances in shaping the radiation dose to organs and tissue near a patient's tumors, leading to effective treatments with less toxicity," added Shireen Parsai, M.D., a radiation oncologist at OhioHealth who led this research during her residency at Cleveland Clinic. "TFRT takes advantage of the differential repair of tumors vs. healthy tissue by modulating how the dose is delivered to nearby tissue over time. This approach gives normal tissue more time to heal, allowing us, in theory, to deliver the same curative doses of [radiation](#) with less detriment to the patient's quality of life."

**More information:** Shireen Parsai et al, In vivo assessment of the safety of standard fractionation Temporally Feathered Radiation Therapy (TFRT) for head and neck squamous cell carcinoma: An R-IDEAL Stage 1/2a first-in-humans/feasibility demonstration of new technology implementation, *Radiotherapy and Oncology* (2021). [DOI: 10.1016/j.radonc.2021.07.023](https://doi.org/10.1016/j.radonc.2021.07.023)

Provided by Cleveland Clinic

Citation: Research proves feasibility of new cancer treatment approach (2021, September 1)  
retrieved 26 April 2024 from

<https://medicalxpress.com/news/2021-09-feasibility-cancer-treatment-approach.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.