

Study links air pollution to nearly 6 million preterm births around the world

September 28 2021



Credit: Unsplash/CC0 Public Domain

Air pollution likely contributed to almost 6 million premature births and almost 3 million underweight babies in 2019, according to a UC San Francisco and University of Washington global burden of disease study

and meta-analysis that quantifies the effects of indoor and outdoor pollution around the world.

The analysis, published September 28, 2021, in *PLOS Medicine*, is the most in-depth look yet at how [air pollution](#) affects several key indicators of pregnancy, including gestational age at birth, reduction in [birth weight](#), [low birth weight](#), and [preterm birth](#). And it is the first [global burden of disease study](#) of these indicators to include the effects of [indoor air pollution](#), mostly from cook stoves, which accounted for two-thirds of the measured effects.

A growing body of evidence points to air pollution as a major cause of preterm birth and low birthweight. Preterm birth is the leading cause of neonatal mortality worldwide, affecting more than 15 million infants every year. Children with low birthweight or who are born premature have higher rates of major illness throughout their lives.

The World Health Organization estimates that more than 90 percent of the world's population lives with polluted outdoor air, and half the [global population](#) is also exposed to indoor air pollution from burning coal, dung and wood inside the home.

"The air pollution-attributable burden is enormous, yet with sufficient effort, it could be largely mitigated," said lead author Rakesh Ghosh, Ph.D., a prevention and public health specialist at the Institute for Global Health Sciences at UCSF.

The analysis, which was conducted with researchers at the Institute for Health Metrics and Evaluation (IHME) at the University of Washington, quantified preterm birth and low birthweight risks based on total indoor and outdoor pollution exposure, while also accounting for the likelihood that the negative effects taper off at higher levels.

The study concluded that the global incidence of preterm birth and low birthweight could be reduced by almost 78 percent if air pollution were minimized in Southeast Asia and sub-Saharan Africa, where indoor pollution is common and preterm birth rates are the highest in the world.

But it also found significant risks from ambient air pollution in more developed parts of the world. In the United States, for example, outdoor air pollution is estimated to have contributed to almost 12,000 preterm births in 2019.

Previously, the same research team [quantified](#) the effects of air pollution on early life mortality, concluding that it contributed to the deaths of 500,000 newborns in 2019.

"With this new, global and more rigorously generated evidence, air pollution should now be considered a major driver of infant morbidity and mortality, not just of chronic adult diseases," Ghosh said. "Our study suggests that taking measures to mitigate climate change and reduce air pollution levels will have significant health co-benefit for newborns."

More information: *PLOS Medicine* (2021).
[journals.plos.org/plosmedicine ... ournal.pmed.07267523](https://journals.plos.org/plosmedicine/...ournal.pmed.07267523)

Provided by University of California, San Francisco

Citation: Study links air pollution to nearly 6 million preterm births around the world (2021, September 28) retrieved 28 April 2024 from <https://medicalxpress.com/news/2021-09-links-air-pollution-million-preterm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.