

Long-term injectable medications plus counseling improve adherence and symptoms in bipolar disorder patients

September 16 2021



Credit: Unsplash/CC0 Public Domain

Researchers have found that adherence to taking medications improved in people with bipolar disorder with the use of long-term injectable meds

combined with psychosocial intervention.

The research results from University Hospitals (UH) Cleveland Medical Center and Case Western Reserve University (CWRU) School of Medicine appear in the September 16, 2021 issue of the journal *The Primary Care Companion for CNS Disorders*.

"Poor [medication adherence](#) is widely prevalent in people with [bipolar disorder](#) and is often associated with negative outcomes," said lead author Martha Sajatovic, MD, Director of the Neurological and Behavioral Outcomes Center, CWRU School of Medicine and UH Cleveland Medical Center. "A growing body of literature focuses on clinical approaches to address poor adherence that may lead to substantial outcome improvements for this group of people."

Bipolar disorder is a mental health condition that causes extreme mood swings that include emotional highs (mania or hypomania) and lows (depression).

The study enrolled 30 people with bipolar disorder who had adherence problems. More than 20 percent of them reported missing their bipolar medication in the past week or month when initially screened.

For the study, the participants were given the long-acting injectable (LAI) antipsychotic medication aripiprazole, approved by the US Food and Drug Administration (FDA) for the treatment of bipolar disorder.

"However, in spite of the adherence advantages for long-acting injectable meds, simply switching individuals to them may not be enough to sustain long-term behavioral change," said Dr. Sajatovic. "Our pilot trial combined the LAI with a brief behavioral approach called customized adherence enhancement (CAE). Together we called them CAE-L to assess the effects on adherence, bipolar disorder symptoms,

and functional status," she said.

CAE was delivered by a trained [social worker](#) following a detailed curriculum and delivered in the same clinical visit when the patients came in for their medication. According to Dr. Sajatovic, barriers to adherence could consist of multiple components, including a lack of education about the medications; communication issues with providers; strategies to enhance medication routines, and substance abuse issues.

The CAE is a brief, practical intervention program designed to address an individual's specific adherence barriers. Each module could be combined with other modules as determined by a screening assessment. For this study, CAE was delivered in seven sessions (an initial baseline session, followed by one each month for the six months of the study).

The medication was also administered once a month for six months. Participants also continued other maintenance treatments such as mood-stabilizing drugs (lithium, valproate, or lamotrigine), or antidepressants, and hypnotic drugs for sleep prescribed for at least one month prior to enrollment.

"Overall, our findings suggest that a personalized intervention to address adherence barriers combined with LAI can significantly improve outcomes in high-risk individuals with bipolar disorder," said Dr. Sajatovic.

At the end of the six months, self-reported adherence behaviors improved. Adherence remained steady and bipolar symptoms improved.

CAE-L was associated with excellent adherence to LAI (100 percent of individuals received injection within one week of the scheduled time), compared with initial screening where individuals missed a mean of 50.1

percent of prescribed oral medication in the past week and 40.6 percent of medication in the past month.

The proportion of missed medications in the past week from screen to 24 weeks significantly improved from 50.1 percent to 16.9 percent, and past month improved from 40.6 percent to 19.2 percent.

From baseline to week 24, there were significant decreases in bipolar disorder symptoms and global psychopathology. Functioning was significantly improved from baseline to week 24. Participants also found CAE intervention was highly acceptable.

The study's limitations include small sample size, no control group for comparison, and all patients coming from a single hospital site. An additional limitation is that adherence was based on self-report, which has potential to undercount missed medication. Twenty-one completed the trial out of the initial 30 who started (nine participants dropped out of the study early for various reasons).

In addition to Dr. Sajatovic, MD, the other authors are Jennifer B. Levin, Ph.D.; Luis F. Ramirez, MD; Kristin A. Cassidy, MA; Nora McNamara, MD; Edna Fuentes-Casiano, MSSA, LSW; Deionte Appling, BA, all with CWRU and UH; Farren B. S. Briggs, Ph.D., with CWRU, and Betsy Wilson, MPH, who was with CWRU and UH at the time of the study.

More information: Martha Sajatovic et al, Long-Acting Injectable Antipsychotic Medication Plus Customized Adherence Enhancement in Poor Adherence Patients With Bipolar Disorder, *The Primary Care Companion For CNS Disorders* (2021). [DOI: 10.4088/PCC.20m02888](https://doi.org/10.4088/PCC.20m02888)

Provided by University Hospitals Cleveland Medical Center

Citation: Long-term injectable medications plus counseling improve adherence and symptoms in bipolar disorder patients (2021, September 16) retrieved 7 May 2024 from <https://medicalxpress.com/news/2021-09-long-term-medications-adherence-symptoms-bipolar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.