

# Outdoor exercise benefits new moms' mental health during the COVID-19 pandemic

September 24 2021, by Iris Lesser, Scott Lear

---



Women who engage in physical activity in the postnatal period report better mental well-being than those who are less physically active. Author provided

The extended duration of the COVID-19 pandemic means more women will give birth during the pandemic, and some will have more than one pregnancy and postpartum experience. As physical activity researchers who advocate for exercise as medicine, we are studying the impact of exercise on well-being of postpartum women during the pandemic.

The COVID-19 pandemic has affected everyone's day-to-day lives, but [mothers](#) have been especially challenged. Women have experienced a negative impact on mental well-being and [physical activity behavior](#), with mothers being especially vulnerable with increased child-care responsibilities. New mothers in the postpartum phase are at an additional disadvantage due the heightened challenge of caring for infants.

## Postpartum challenges

The [postpartum phase](#) is often defined as the first six weeks after childbirth when post-pregnancy [physical changes](#) such as uterine shrinking and hormonal fluctuations are the greatest. These changes can also greatly affect the mental health of new mothers—the prevalence of depression is approximately 15 percent. However, it is possible for new mothers to continue to experience post-partum effects for up to one year.

Since the onset of COVID-19, mental health issues have increased among postpartum [women](#), with [41 percent reporting depression and 72 percent reporting moderate-to-high anxiety](#) (compared to 15 percent and 29 percent pre-pandemic).

Pre-pandemic challenges such as sleep deprivation, lack of self-care or medically complicated deliveries are now exacerbated. For example, [women who gave birth during the pandemic](#) may have experienced reduced direct maternal care, a lack of a home support system due to visiting restrictions and a lack of in-person breastfeeding support.

Physical activity is an effective therapy for anxiety and depression and may be beneficial for new mothers. As little as a single group [exercise session of 45 minutes can improve anxiety levels](#) in women without a history of mental illness. Women who engage in physical activity in the

postnatal period [report better mental well-being](#) than those who are less physically active.

However, physical activity levels are [lower in the postpartum period](#) than prior to pregnancy. This is despite the American College of Obstetrics and Gynecologists recommending that women slowly [return to physical activity four to six weeks after giving birth](#) and work their way up to the general physical activity guidelines of 150 minutes of moderate to vigorous physical activity per week. [Barriers in the postnatal period](#) include reduced motivation due to a lack of social support and childcare responsibilities.

## **Pandemic-safe exercise solutions**

At 18 months into the pandemic with no end in sight, and safe indoor opportunities for [exercise](#) less accessible, [moms are in need of innovation](#). An [outdoor environment](#) has the advantage of being easily accessible and COVID-19 safe, and having additional beneficial effects. For instance, exercising in nature, known as "[green exercise](#)" has been associated with reductions in anxiety and stress, while being outside has been shown to increase enjoyment of exercise.

With the high degree of [safety in an outdoor environment](#), could we support new moms with safe physical activity programming outdoors? We were interested in providing moms with the tools they need to return to physical activity safely and ensure they have the added competence to engage in physical activity with a new baby. If we want moms to thrive in this new normal we need to help them find their way back to movement and optimal mental well being.

Working with a team at the University of the Fraser Valley, in the spring of 2021 we enrolled 21 women who were less than nine months postpartum in an eight-week, biweekly outdoor group exercise program.

As a mother of two (including one born in the midst of the pandemic), Iris (one of the co-authors of this article) had both personal and professional interest in finding ways to meet the physical activity needs of new mothers.

Engaging in the program in an outdoor location with physical distancing allowed women to fully participate without risk of exposure to COVID-19. Mothers reported a decrease in depression after the program, along with an improvement in motivation and in meeting basic psychological needs. Perceived stress and anxiety also were reduced, but not significantly.

## Ongoing research

These findings suggest that new mothers may experience an improvement in overall well-being after engaging in group exercise. This form of group exercise addressed a number of the common barriers to exercise participation during the postnatal phase. This includes the opportunity to bring their child to the program, engaging with other new mothers and the program being designed specifically for postpartum fitness. To determine the long-term effects on physical activity, we plan to conduct a followup at six months.

Supporting postpartum women with physical [activity](#) opportunities may help them exercise more in the postpartum period. This may further improve mental well-being in this at-risk population.

However, there is still a large gap in the literature regarding women's postnatal [physical activity](#) experiences. We are currently recruiting women less than 12 months postpartum to [complete an online survey](#) to help us learn more.

This article is republished from [The Conversation](#) under a Creative

Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Outdoor exercise benefits new moms' mental health during the COVID-19 pandemic (2021, September 24) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-09-outdoor-benefits-moms-mental-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.