

Oxcarbazepine tied to nontrauma fracture in children aged 4 to 13 with epilepsy

September 22 2021



(HealthDay)—For children aged 4 to 13 years with epilepsy, initiating



oxcarbazepine, but not levetiracetam, antiseizure medication (ASM) is associated with an increased risk for nontrauma fracture, according to a study published in the September issue of *Epilepsia*.

Daniel G. Whitney, Ph.D., from the University of Michigan in Ann Arbor, and colleagues extracted claims data for <u>children</u> aged 4 to 13 years with at least five years of continuous health plan enrollment. Children with epilepsy who were ASM-naive were grouped according to treatment initiation, including <u>levetiracetam</u> or oxcarbazepine, and were compared to children without epilepsy and without ASM exposure. For up to four years of follow-up, the crude incidence rate (IR, per 1,000 person-years) and IR ratio (IRR) were estimated for nontrauma fracture.

The researchers found that the crude IR for nontrauma fracture was 21.5, 19.8, and 34.4 for non-ASM users without epilepsy, those taking levetiracetam, and those taking oxcarbazepine, respectively. The crude IRR of nontrauma fractures was similar for levetiracetam and non-ASM users (IRR, 0.92; 95 percent confidence interval, 0.63 to 1.34) and was elevated for oxcarbazepine (IRR, 1.60; 95 percent confidence interval, 1.09 to 2.35); compared with levetiracetam, the crude IRR was elevated for oxcarbazepine (IRR, 1.74; 95 percent confidence interval, 1.02 to 2.99). After adjustment for covariates, the findings were consistent, except for the comparison of oxcarbazepine with levetiracetam (hazard ratio, 1.71; 95 percent confidence interval, 0.99 to 2.93).

"Knowing that oxcarbazepine is associated with a higher fracture risk will allow clinicians to be more thoughtful about to which children they prescribe the medication," a coauthor said in a statement.

More information: <u>Abstract/Full Text (subscription or payment may be required)</u>



Copyright © 2021 HealthDay. All rights reserved.

Citation: Oxcarbazepine tied to nontrauma fracture in children aged 4 to 13 with epilepsy (2021, September 22) retrieved 17 July 2024 from

https://medicalxpress.com/news/2021-09-oxcarbazepine-tied-nontrauma-fracture-children.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.