

## New potential factor contributing to severity of COVID-19 identified

September 24 2021



Credit: Pixabay/CC0 Public Domain

The University of Kent's School of Biosciences and the Institute of Medical Virology at Goethe-University, Frankfurt am Main, have identified a protein that may critically contribute to severe forms of COVID-19.

SARS-CoV-2 is the coronavirus that causes COVID-19. While many



individuals develop only mild or no symptoms upon SARS-CoV-2 infection, others develop severe, life-threatening disease.

Researchers have found that the infection of cells with SARS-CoV-2 results in increased levels of a protein called CD47 on the cell surface.

CD47 is a so-called 'do not eat me' signal to the immune system's defenses that protect cells from being destroyed. Virus-induced CD47 on the surface of infected cells is likely to protect them from immune system recognition, enabling the production of larger amounts of virus, resulting in more severe disease.

Well-known <u>risk factors</u> for severe COVID-19 such as older age and diabetes are associated with higher CD47 levels. High CD47 levels also contribute to <u>high blood pressure</u>, which is a large risk factor for COVID-19 complications such as heart attack, stroke, and kidney disease.

The data suggest that age and virus-induced high CD47 levels contribute to severe COVID-19 by preventing an effective immune response and increasing disease-associated tissue and organ damage.

Since therapeutics targeting CD47 are in development, this discovery may result in improved COVID-19 therapies.

Professor Martin Michaelis, University of Kent, says that "this is exciting. We may have identified a major factor associated with severe COVID-19. This is a huge step in combatting the disease and we can now look forward to further progress in the design of therapeutics."

Professor Jindrich Cinatl, Goethe-University Frankfurt, says that "these additional insights into the disease processes underlying COVID-19 may help us to design better therapies, as well as appreciation for the



importance of the breadth of research being conducted. Through this avenue, we have achieved a major breakthrough and exemplified that the fight against the disease continues."

The study is published by the <u>scientific journal</u> Current Issues in Molecular Biology.

**More information:** Katie-May McLaughlin et al, A Potential Role of the CD47/SIRPalpha Axis in COVID-19 Pathogenesis, *Current Issues in Molecular Biology* (2021). DOI: 10.3390/cimb43030086

Provided by University of Kent

Citation: New potential factor contributing to severity of COVID-19 identified (2021, September 24) retrieved 6 June 2024 from <u>https://medicalxpress.com/news/2021-09-potential-factor-contributing-severity-covid-.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.