

New study reveals signs of dementia are written in the blood

September 13 2021





PCA of 33 AD or 6 selected dementia-related compounds showed significant differences between patients with dementia and HE subjects. Blood data of dementia and HE subjects were subjected to PCA. (A) A PCA plot using abundances of 33 (7 increased and 26 decreased) dementia-related compounds.



Dementia and HE subjects were separated into two domains (see text). Red, dementia; green, HE. (B) PCA was also performed using only 6 selected dementia markers (dimethyl-guanosine, pseudouridine, S-methyl-ET, ET, trimethyl-histidine, and NADP+). Credit: DOI: 10.1073/pnas.2022857118

Scientists in Japan have identified metabolic compounds within the blood that are associated with dementia.

The study revealed that the levels of 33 metabolites differed in patients with <u>dementia</u>, compared to <u>elderly people</u> with no existing health conditions. Their findings, published this week in *PNAS*, could one day aid diagnosis and treatment of dementia.

"Metabolites are <u>chemical substances</u> produced by vital chemical reactions that occur within <u>cells</u> and tissues," said first author Dr. Takayuki Teruya, who works in the G0 Cell Unit at the Okinawa Institute of Science and Technology Graduate University (OIST). "Our body normally keeps these levels in balance, but as we age and if we develop diseases like dementia, these levels can fluctuate and change."

Dementia is not just a single disease, but a general term used to describe a set of symptoms, including a slow but typically irreversible decline in the ability to remember, think, make decisions or perform day-to-day activities. Of all aging-associated diseases, dementia is one of the most serious, not only for the patients and their family but for society as a whole, with an estimated 55 million people living with the disease worldwide.

While scientists know that dementia is caused by damage to nerves, the exact cause of this damage, and methods as to how it can be detected and treated have remained elusive.



In the study, the research team analyzed samples of blood collected from eight patients with dementia, as well as eight healthy elderly people. They also collected samples from eight healthy young people to use as a reference. Unlike most studies analyzing <u>blood metabolites</u>, this research included compounds found within red blood cells.

"Blood cells are difficult to handle because they undergo metabolic changes if left untreated even for a short period of time," explained Dr. Teruya.

However, the research team recently developed a way to stabilize metabolites in red blood cells, allowing them to examine for the first time the relationship between red blood cell activity and dementia.



33 dementia markers			Dementia							HE								
			#1	#2	#3	#4	#5	#6	#7	#8	#9	#10	#11	#12	#13	#14	#15	#16
Higher in dementia (7)	A	Quinolinic acid	52.4	59.9	56.2	48.0	51.6	68.9	76.9	45.6	38.4	53.8	39.4	36.8	48.2	46.0	40.6	64.6
		Dimethyl-guanosine	49.2	63.0	51.8	42.3	60.9	79.7	58.3	62.2	42.7	49.8	52.7	38.4	52.8	47.9	50.8	47.0
		Pseudouridine	53.3	56.8	54.1	45.3	55.7	84.4	65.2	54.8	41.5	54.3	50.0	38.8	53.9	49.2	49.2	49.4
		Indoxyl-sulfate	57.1	63.4	53.2	53.6	43.7	78.1	58.8	57.2	37.3	59.9	47.4	38.1	52.8	52.4	39.1	38.7
		Kynurenine	67.0	50.9	56.7	37.6	63.3	53.5	57.2	60.1	47.9	52.6	50.9	22.8	44.4	55.3	44.4	51.5
		N6-Acetyl-lysine	85.1	46.3	45.4	49.7	57.8	48.4	71.7	44.9	45.8	48.2	42.9	44.5	47.8	44.8	43.8	46.0
		Adenosine	69.8	51.9	43.1	44.0	50.3	50.9	45.6	51.5	43.0	45.8	43.4	46.3	43.1	50.4	40.2	39.5
Lower in dementia (26)	в	S-Methyl-ergothioneine	40.7	40.2	44.2	40.5	41.2	38.9	40.2	42.0	52.5	45.2	57.6	70.9	45.2	57.7	79.0	45.5
		Ergothioneine	40.5	42.8	41.8	43.4	42.0	41.5	37.3	49,9	52.0	44.6	53.1	71.0	49.9	68.0	77.8	45.3
		Trimethyl-histidine	38.9	48.4	51.5	44.1	42.2	43.6	37.4	44.6	65.5	54.0	46.3	69.2	64.4	55.1	77.5	45.6
		Trimethyl-tryptophan	41.1	43.0	42.0	43.0	42.0	41.2	46.3	43.2	56.6	76.3	44.5	59.8	67.6	65.3	51.8	47.2
		Trimethyl-phenylalanine	46.9	46.9	46.9	46.9	46.9	46.9	46.9	47.0	47.2	46.9	47.2	78.2	86.2	49.0	47.1	46.9
		Trimethyl-tyrosine	45.5	45.7	45.7	45.8	45.7	45.5	45.5	45.5	45.6	45.9	47.4	46.9	93.6	47.4	57.5	45.9
	c	Pantothenate	42.0	41.6	40.1	39.1	43.9	38.6	59.7	50.1	69.2	77.2	45.5	51.3	57.7	47.7	52.5	45.6
		Gluconate	43.8	44.4	64.1	33.6	42.5	32.2	56.1	39.8	50.9	53.4	49.5	51.5	76.4	59.1	52.1	60.5
		S-Adenosyl-methionine	42.5	41.3	41.2	41.3	60.6	42.1	43.9	43.6	53.7	46.4	45.8	51.0	44.7	53.1	43,9	45.9
		NADP+	35.1	40.4	33.6	31.6	48.0	42.2	47.2	44.8	55.5	45.2	57.7	58.5	46.8	45.4	53.7	45.3
		Glutathione disulfide	37.3	51.5	22.0	42.0	46.0	46.5	36.2	56.3	62.3	58.7	61.2	58.5	47.2	50.1	58.4	44.0
		ATP	40.8	45.4	30.9	31.9	48.0	46.3	47.1	51.0	42.3	51.5	60.2	60.7	58.5	50.8	61.9	51.0
	D	Methionine	35.3	52.9	40.2	39.8	41.2	39.5	30.7	69.3	54.1	52.1	65.4	56.4	51.4	48.0	50.5	53.5
		Tryptophan	36.9	53.9	43.6	44.7	42.1	40.7	22.5	58.4	52.5	52.9	56.8	49.3	44.7	61.4	61.9	49.8
		Glutamine	44.6	45.2	33,1	40.6	41.2	53.6	33.6	66.9	49.3	51.3	54.8	54.9	60.9	54.9	52.6	67.6
		Betaine	45.4	52.5	44.4	45.8	40.7	42.6	27.3	52.4	52.8	59.5	53.1	56.0	62.3	46.3	60.8	51.3
		Phenylalanine	36.9	59.2	41.6	41.4	37.8	48.5	41.7	58.3	47.2	59.3	69.5	47.6	47.5	59.4	58.8	58.5
		Tyrosine	41.3	38.6	39.6	44.9	44.7	40.5	39.3	59.7	44.5	67.3	60.5	53.5	52.9	56.4	53.7	51.2
		Histidine	44.4	49.4	48.4	49.0	41.8	33.9	30.6	55.6	43.5	64.7	58.4	42.7	76.6	65.2	53.9	52.5
		Uridine	42.9	43.0	45.7	42.9	37.6	33.8	34.2	55.9	53.9	48.2	49.1	49.1	59.8	68.3	62.7	51.5
		Keto(iso)leucine	40.8	46.1	46.1	43.2	31.7	51.8	47.8	46.0	56.2	56.3	65.3	41.1	50.3	69.4	56.8	50.0
		Glycerophosphocholine	48.8	48.8	43.4	35.5	43.6	49.5	37.8	43.3	52.0	50.5	45.6	46.2	55.7	53.5	47.0	50.6
		2-Hydroxybutyrate	48.3	46.9	51.9	40.9	34.7	47.0	66.3	48.6	48.8	41.3	74.7	61.5	56.0	57.0	53.2	53.6
		Dodecanoyl-carnitine	39.5	40.0	39.5	42.0	44.7	55.5	55.1	39.7	54.5	51.0	59.6	44.9	40.0	46.6	60.3	62.2
	E	Caffeine	44.8	43.5	43.6	44.4	43.4	43.7	44,7	45.2	45.7	54.3	55.5	43.3	62.7	66.8	49.2	45.6
		Dimethyl-xanthine	57.7	40.4	40.7	40.7	40.1	40.4	40.3	42.3	45.5	60.7	59.2	39.8	64.2	71.1	55.6	44.3

Red shows high levels of a compound, blue shows low levels of a compound. Heat map reveals the link between certain metabolites and dementia. Compounds in sub-group A were typically higher in dementia patients and lower in healthy elderly people. Compounds in sub-group B-E showed the opposite effect. Credit: Okinawa Institute of Science and Technology

The scientists measured the levels of 124 different metabolites in whole blood and found that 33 metabolites, split into 5 different sub-groups, correlated with dementia. Seven of these compounds increased in dementia patients, whilst 26 of these compounds showed a decrease in



levels. 20, including nine that were abundant in red blood cells, of these compounds had not previously been linked to dementia.

"Identification of these compounds means that we are one step closer to being able to molecularly diagnose dementia," said senior author of the study, Professor Mitsuhiro Yanagida, who leads the G0 Cell Unit at OIST.

The seven metabolites that showed increased levels in patients with dementia were found within the <u>blood</u> plasma and belonged to sub-group A of metabolites. Importantly, some of these compounds are believed to have toxic effects on the central nervous system.

"It's still too early to say, but it could suggest a possible mechanistic cause of dementia as these compounds may lead to impairment of the brain," said Prof. Yanagida.

The research team plans to test this idea in the next steps of their research, by seeing if increases in these metabolites can induce dementia in animal models, like mice.

The remaining 26 compounds that decreased in patients with dementia, compared to healthy elderly people, belonged to four other <u>metabolite</u> sub-groups, B-E.

Six metabolites that decreased in dementia patients were classified into sub-group B, due to their similar structure. These metabolic compounds are antioxidants, which protect cells and tissues by reducing damage caused by free radicals—unstable molecules produced by chemical reactions in cells. The researchers found that these antioxidant compounds derived from food were highly abundant in red <u>blood cells</u> of healthy elderly people.



"It could be that <u>red blood cells</u> deliver not only oxygen but also crucial metabolites that protect the nervous system from damage," said Dr. Teruya.

The remaining sub-groups contain compounds that the researchers believe play a role in supplying nutrients, maintaining energy reserves and protecting neurons from damage.

"In the future, we hope to start some intervention studies, either by supplementing dementia patients with metabolic <u>compounds</u> in subgroups B-E, or by inhibiting the neurotoxins from sub-group A, to see if that can slow, prevent, or even reverse symptoms of dementia," said Prof. Yanagida.

More information: Takayuki Teruya et al, Whole-blood metabolomics of dementia patients reveal classes of disease-linked metabolites. *PNAS* (2021). DOI: 10.1073/pnas.2022857118

Provided by Okinawa Institute of Science and Technology

Citation: New study reveals signs of dementia are written in the blood (2021, September 13) retrieved 25 June 2024 from <u>https://medicalxpress.com/news/2021-09-reveals-dementia-written-blood.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.