

SARS-CoV-2 transmission model suggests primary school infection could be greater this autumn than in 2020

September 14 2021



Credit: Unsplash/CC0 Public Domain

The SARS-CoV-2 epidemic has already had a major impact on children's education, with schools having been required to implement



infection control measures that have led to long periods of absence and classroom closures. With the new school year underway, risk modeling specialists at the University of Bristol have developed a new epidemiological model for SARS-CoV-2 transmission that forecasts primary school infection outbreaks could be more frequent and possibly substantially larger this autumn than in 2020, due to the more transmissive and infectious Delta variant and projected increase in community infection.

The study, published on the preprint server *medRxiv*, is led by Dr. Mark Woodhouse from Bristol's Schools of Earth Sciences and Mathematics, working with colleagues in Earth Sciences, Bristol Veterinary School and with specialists in public health and infectious diseases on the CoMMinS (COVID-19 Mapping and Mitigation in Schools) project.

The new agent-based model characterizes the numbers of daily close contacts that can occur in primary school classrooms, involving pupils and teachers and provides a basis for assessing likely rates of COVID-19 infection transmission and infections within classrooms in the new term. The model takes into account the infection dynamics of the new Delta variant of SARS-CoV-2 and changing guidance for mitigation measures in schools.

The pre-print reports advances in computational probabilistic modeling for estimating in-school infection risks, building on earlier studies by Bristol colleagues (Sparks et al; Aspinall et al). The new analytical framework is based on developments in characterizing, statistically, the complexities of child-with-child and child-with-teacher interactions within the classroom, and how these are modulated by classroom management controls for reducing infection spread.

A previous study (Sparks et al) demonstrated that pupils' interactions with one another have a distinctive statistical profile: some children are



not extensive mixers, whilst others are very sociable and have many, many contacts in the school day. The latter children may act as "superspreaders" in a class or school if they pick up a COVID-19 infection in the community or from another child.

Mitigation strategies to control infection are modeled to allow analysis of their effectiveness in supressing infection outbreaks in classrooms and in limiting pupil absence. The model was applied to re-examine COVID-19 in schools in the UK in autumn 2020 and, with some fine-tuning, is now run to forecast infection levels in autumn 2021 when the more infectious Delta variant is anticipated to be dominant and school transmission could be a major concern in any new wave of the epidemic.

Using forecasts that are tracking current upward trends in incidence rates in the UK, key findings and implications from the model include:

- Substantial COVID-19 outbreaks could occur in primary schools during the Autumn term.
- A small proportion of primary schools (less than one percent) might experience outbreaks, as defined by the DfE1 and that there will be more such outbreaks than in 2020.
- Most primary schools are expected to experience at least one classroom infection case.
- A comparison of alternative mitigation measures indicates that bubble quarantining is less effective in reducing infection in primary schools when compared to testing-based surveillance and removing children when they become symptomatic.
- Bubble quarantining is highly disruptive to education but offers little public health benefit. However, bubble quarantining may be considered necessary to manage large outbreaks in primary schools1.
- Lateral flow testing-based surveillance achieves a greater reduction in numbers of infected pupils compared to bubble



quarantining. Confirmatory PCR testing has very little benefit in reducing infection levels in the classroom but can reduce quarantine time for false-positive rapid tests.

• The study confirms that individual school's policies that encourage social distancing together with other measures like ventilation and hygiene and routine lateral flow testing can reduce infection transmission, but the researchers recognize there are practical limits to such policies, especially with primary school age children.

The study's findings indicate that testing-based surveillance of infections in the classroom population with isolation of positive cases is a more effective mitigation measure than bubble quarantining both for reducing transmission in schools and for avoiding pupil absence, even accounting for insensitivity of self-administered tests. Bubble quarantining results in large numbers of pupils absent from school, with only modest impact of classroom infection. However, maintaining a reduced contact rate within the classroom has a major beneficial impact for managing COVID-19 in school settings.

Dr. Mark Woodhouse, Knowledge Exchange Fellow in the Schools of Earth Sciences and Mathematics, said: "Schools re-opening for the new school year face the prospect of managing and mitigating risks associated with the evolving COVID-19 pandemic outbreak in England and, in particular, the aggressive dynamics of the Delta variant of the coronavirus, which is growing in prevalence.

"Mathematical models are being used extensively to help understand the probability of the virus being transmitted between persons, to inform prevention and control strategies. For schools, our findings suggest there are substantive implications for the relative efficacies of different mitigation measures that may be followed. For instance, our modeling presents evidence that, with the emergence of the Delta variant and other



changing factors, there is likely to be a distinct contrast between the efficacies of bubbles and of regular testing when it to comes to projected numbers of infections in schools and pupil absences in the first few weeks of the new academic year. In addition, our results indicate potential benefits from refocusing on social distancing in schools, as well as enhanced ventilation and good hygiene."

Professor Sir Stephen Sparks FRS, Emeritus Professor in the School of Earth Sciences at the University of Bristol, added: "We believe our study projections provide important information for policy considerations, especially in relation to absences from school that will be consequent on choices of different mitigation measures. There is the prospect that schools could become driver reservoirs of community infection, rather than the other way around, as was the case with the Alpha variant.

"These, and related arguments from our modeling, suggest that—in forecast mode—our stochastic contact network framework should be of value to Directors of Public Health and Education Authorities charged with responsibility for prompt assessment and managing of infection transmission dynamics in classes, schools and in multiple schools in a given education authority area or wider region. In terms of current or future DfE criteria for school actions, our model can be tailored straightforwardly to different schools' demographics and to the relevant local community incidence level."

More information: M. J. Woodhouse et al, Analysis of alternative COVID-19 mitigation measures in school classrooms: an agent-based model of SARS-CoV-2 transmission, (2021). <u>DOI:</u> <u>10.1101/2021.08.30.21262826</u>

R. S. J. Sparks et al, A novel approach for evaluating contact patterns and risk mitigation strategies for COVID-19 in English primary schools with application of structured expert judgement, *Royal Society Open*



Science (2021). DOI: 10.1098/rsos.201566

W.P. Aspinall et al, Quantifying threat from COVID-19 infection hazard in Primary Schools in England, (2020). DOI: 10.1101/2020.08.07.20170035

Provided by University of Bristol

Citation: SARS-CoV-2 transmission model suggests primary school infection could be greater this autumn than in 2020 (2021, September 14) retrieved 6 May 2024 from https://medicalxpress.com/news/2021-09-sars-cov-transmission-primary-school-infection.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.