

Women with heart problems are treated differently than men

September 20 2021



Credit: Pin Adventure Map/Unsplash

Women with some cardiac conditions receive less evidence-based treatment than their male counterparts, research published today by the *Medical Journal of Australia* has confirmed.

Researchers from the University of Sydney, Westmead Hospital and Concord Repatriation General Hospital, assessed differences in the evidence-based treatment received by men and women with non-ST-elevation [acute coronary syndromes](#) (NSTEMI) and in their outcomes, both in-hospital and at 6-month follow-up. They also separately assessed these differences in patients with documented coronary artery disease (CAD).

"The proportion of women who underwent cardiac catheterisation was smaller (1710, 71 percent v 4134, 77 percent), and the median time to catheterisation was longer (53 h; IQR, 28–91 h v 47 h; IQR, 25–77 h) than for men; non-obstructive [coronary artery disease](#) (NOCAD) was detected in a larger proportion of women than men during catheterisation (602, 35 percent v 566, 14 percent)," they authors, led by Professor David Brieger, Professor of Cardiology at the University of Sydney.

"At discharge, fewer women were prescribed aspirin (85 percent v 91 percent), a second antiplatelet medication (59 percent v 68 percent), beta-blockers (71 percent v 75 percent), or statins (86 percent v 92 percent), or referred to cardiac rehabilitation (54 percent v 63 percent).

"A total of 4676 patients had documented CAD, including 1108 women (24 percent). Smaller proportions of women with CAD than of men underwent [coronary artery bypass](#) grafting (110, 10 percent v 563, 16 percent) or were prescribed statins at discharge (94 percent v 96 percent).

"Fewer women than men were referred to cardiac rehabilitation (750, 69 percent v 2652, 75 percent), including of those who had been revascularised (CABG: 97, 77 percent v 509, 83 percent; PCI: 480, 76 percent v 1623, 81 percent)," Brieger and colleagues reported.

"The larger proportion of women with NOCAD may partly explain the difference. However, NOCAD is not a benign condition, and patients can benefit from secondary prevention therapies," they wrote.

"In Australia, adherence to guideline-based therapy for people with NSTEMI could be improved, especially for [women](#) in hospital and for both sexes at discharge."

More information: Bianca C Bachelet et al, Sex differences in the management and outcomes of non-ST-elevation acute coronary syndromes, *Medical Journal of Australia* (2021). [DOI: 10.5694/mja2.51220](#)

Provided by University of Sydney

Citation: Women with heart problems are treated differently than men (2021, September 20) retrieved 26 April 2024 from <https://medicalxpress.com/news/2021-09-women-heart-problems-differently-men.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--