

# Data shows Americans need at least 5 hours per week of physical activity to prevent some cancers

October 14 2021

---



Credit: CC0 Public Domain

A new report finds more than 46,000 cancer cases annually in the United States could be prevented if Americans met the 5 hours per week of

moderate-intensity recommended physical activity guidelines. The latest data appearing in the journal *Medicine & Science in Sports & Exercise* show 3% of all cancer cases in adults in the U.S. aged 30 years and older during 2013 to 2016 were attributable to physical inactivity and the proportion was higher in women (average annual attributable cases 32,089) compared to men (14,277).

For both men and women, states with the highest proportion of cancers attributable to physical inactivity were in the South, such as Kentucky, West Virginia, Louisiana, Tennessee, and Mississippi, whereas the lowest proportions were found in the Mountain region and northern states, such as Utah, Montana, Wyoming, Washington, and Wisconsin.

Led by Adair Minihan, MPH at the American Cancer Society, this is the first study to estimate the number of cancer cases attributable to physical inactivity based on cancer sites (breast, endometrial, colon, stomach, kidney, esophageal adenocarcinoma, and urinary bladder) by state. Data show when focusing on specific cancer sites, 16.9% of stomach cancers, 11.9% of endometrial cancers, 11.0% of kidney cancers, 9.3% of colon cancers, 8.1% of esophageal cancers, 6.5% of female breast cancers, and 3.9% of urinary bladder cancers were associated with lack of exercise. By state, the proportion of cancer cases attributable to [physical inactivity](#) ranged from 2.3% in Utah to 3.7% in Kentucky.

While this data shows the importance of physical activity, there are many barriers to recreational physical activity, which include, but are not limited to, lack of time due to long working hours in low-wage jobs, the cost of gym memberships or personal equipment, lack of access to a [safe environment](#) in which to be active, and potential childcare costs involved with recreational physical activity. Unfortunately, these barriers are more likely to affect historically marginalized populations, including the Black population and individuals with a limited income, underscoring the importance of enhancing health equity.

"These findings underscore the need to encourage physical activity as a means of [cancer](#) prevention and implement individual- and community-level interventions that address the various behavioral and socioeconomic barriers to recreational physical activity," write the authors. "Understanding and reducing the behavioral and socioeconomic barriers to physical activity is essential for optimizing intervention strategies targeting at risk groups across the country."

**More information:** Adair K. Minihan et al, Proportion of Cancer Cases Attributable to Physical Inactivity by US State, 2013-2016, *Medicine & Science in Sports & Exercise* (2021). [DOI: 10.1249/MSS.0000000000002801](#)

Provided by American Cancer Society

Citation: Data shows Americans need at least 5 hours per week of physical activity to prevent some cancers (2021, October 14) retrieved 6 May 2024 from <https://medicalxpress.com/news/2021-10-americans-hours-week-physical-cancers.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--