

# A breakdown in communication: Mitochondria of diabetic patients can't keep time

October 20 2021

---



Mitochondria. Credit: Wikipedia commons

Almost all cells regulate their biological processes over a 24-hour period, otherwise called a cell's circadian rhythm. To do so, cells use a biological clock that cycles different genes on and off throughout the day and

night. Scientists already know that our metabolic health can suffer when our biological clock breaks down, due to shift work or sleep disorders, for example. However, it's unclear how exactly the biological clock of people with type 2 diabetes differs from that of healthy people.

Now a team of international scientists has shown that the skeletal muscle in people with type 2 [diabetes](#) has a different circadian rhythm. They argue that this might arise because of a communication breakdown between a cell's timekeeping molecules and mitochondria, which produce chemical energy for [cells](#).

"The promise of this research is that it may help us to fine-tune the timing of interventions and other medications to treat type 2 diabetes, in order to optimize their effectiveness," says Professor Juleen R. Zierath from Karolinska Institutet and the Novo Nordisk Foundation Center for Basic Metabolic Research (CBMR) at the University of Copenhagen.

## **A different pattern of daily gene expression**

In the study, which was published in *Science Advances*, the scientists first obtained [skeletal muscle](#) cells from people with type 2 diabetes and measured which genes showed cycling behavior over two days and compared them with cells from similar healthy people. They discovered that cells from people with type 2 diabetes had fewer—and some different—cycling genes.

They carried out further experiments using data generated from clinical tests in people with type 2 diabetes and mice, as well as cell-based experiments. These experiments demonstrated that mitochondria communicate with the molecules that keep time in our cells, and that this communication is disrupted in people with type 2 diabetes.

## Diabetes treatments may be more effective if timed to the body clock

Some of the most widely used pharmacological treatments for type 2 diabetes affect mitochondria, meaning that they may work differently depending on the time of day they are taken. As a result, these findings highlight the importance of considering cellular rhythms when prescribing treatments for type 2 diabetes.

"Exercise and diet are regularly used treatment interventions for people with type 2 diabetes, and both of these treatments can affect the time-keeping molecules and mitochondria," says Dr. Brendan Gabriel from the Department of Physiology and Pharmacology at Karolinska Institutet.

Brendan Gabriel is first author on the paper together with Assistant Professor Ali Altintas from CBMR.

"Given that disrupted sleeping patterns are known to be associated with an [increased risk](#) of developing type 2 diabetes, our findings provide evidence of how these disruptions may link to the molecular biology within cells," says Ali Altintas.

**More information:** Brendan M. Gabriel et al, Disrupted circadian oscillations in type 2 diabetes are linked to altered rhythmic mitochondrial metabolism in skeletal muscle, *Science Advances* (2021).

[DOI: 10.1126/sciadv.abi9654](https://doi.org/10.1126/sciadv.abi9654).

[www.science.org/doi/10.1126/sciadv.abi9654](https://www.science.org/doi/10.1126/sciadv.abi9654)

Provided by University of Copenhagen

Citation: A breakdown in communication: Mitochondria of diabetic patients can't keep time (2021, October 20) retrieved 20 April 2024 from <https://medicalxpress.com/news/2021-10-breakdown-mitochondria-diabetic-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.