


Clot-busting drug may be safe for people with stroke and unruptured aneurysms

October 6 2021



Credit: CC0 Public Domain

Clot-busting drugs are the main treatment for stroke, but they have often not been recommended for people who have an aneurysm in their brains that has not ruptured. An aneurysm is a bulge in the wall of a blood vessel in the brain. A new study finds that these aneurysms rarely burst after treatment with clot-busting drugs, so the drugs may be safe for use. The study is published in the October 6, 2021, online issue of *Neurology*.

"These results are exciting because they may expand the population who can receive these thrombolytic therapies that can prevent death and disability for people with strokes," said study author Jyri Juhani Virta, MD, Ph.D., of the University of Helsinki in Finland.

For the study, researchers looked at the records for all people with a [stroke](#) who were treated with clot-busting drugs at a stroke center in Finland over 15 years. After determining how many had unruptured aneurysms, they then looked to see how many people had a hemorrhagic stroke, or an [intracranial hemorrhage](#), caused by the rupture of the [aneurysm](#) while they were in the hospital after the treatment.

Of the 3,953 people in the study, 132 people had one or more unruptured aneurysms. Of the 155 unruptured aneurysms, 141 were the type called saccular aneurysms and 14 were fusiform aneurysms.

The aneurysms ruptured after treatment causing a [hemorrhagic stroke](#) in only three people, or 2.3%. All three had large aneurysms called fusiform basilar artery aneurysms. All three were treated with blood thinners after their treatment. None of the saccular type of aneurysms ruptured.

"Fortunately, today blood thinners are not commonly used early after [ischemic stroke](#), so that may help to reduce this risk even further," Virta said. "Our results suggest that thrombolytic therapy may be relatively safe for people with ischemic stroke even if they have larger saccular aneurysms that have not ruptured. For people with large fusiform aneurysms, additional consideration should be taken regarding treatment as [blood thinners](#) may increase the risk of aneurysm rupture."

A limitation of the study was that only a small number of people had large aneurysms.

Provided by American Academy of Neurology

Citation: Clot-busting drug may be safe for people with stroke and unruptured aneurysms (2021, October 6) retrieved 10 May 2024 from <https://medicalxpress.com/news/2021-10-clot-busting-drug-safe-people-unruptured.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.