

Common infections in pregnant mums linked to leukemia risk in children

October 28 2021



Credit: Unsplash/CC0 Public Domain

Common maternal infections during pregnancy, such as urinary and

respiratory tract infections were linked to a child developing leukemia, a new study has found. But the increase in overall risk was small given the rarity of childhood leukemia.

Researchers looked at whether childhood leukaemias including acute lymphoblastic leukemia and acute myeloid leukemia or any other childhood [cancer](#) occurred more frequently in children whose mothers had self-reported common infections during their pregnancy.

The study, led by the Murdoch Children's Research Institute (MCRI) and the University of Oxford and published in the *International Journal of Epidemiology*, found that while [urinary tract](#) and [respiratory tract infections](#) were associated with childhood leukemia, the increase in overall risk was small given the rarity of a child developing this form of cancer.

The International Childhood Cancer Cohort Consortium (I4C) study analyzed data from Australia, Denmark, Israel, Norway, the UK and the US. More than 300,000 children were recruited for the study over 60 years between the 1950s until the 2000s.

Senior author of the study and Director of I4C International Data Coordinating Centre, based MCRI, Professor Terry Dwyer, said that the results were reassuring for parents.

"Previous studies have found a link between maternal infections and childhood leukemia, however evidence from large cohort studies tracking children over time, was lacking until this research," he said.

Professor Dwyer said innovative projects like the I4C study could provide the greatest opportunity to find a preventable cause of childhood cancer.

MCRI Professor and GenV Scientific Director Melissa Wake said the results of this study demonstrated the immense value of large, whole-of-population cohort studies, such as MCRI's GenV research project.

"This I4C study is the largest combined prospective study of factors that might increase childhood leukemia risk to date. Only by looking at data from such a large cohort were researchers able to confidently conclude that while leukaemias are the most common [childhood](#) cancer, they are still rare," she said.

"Similarly, GenV aims to be the world's largest birth and parent [cohort](#) and has the potential to provide researchers with child and adult health and wellbeing data on a similar scale in the future. Every family in Victoria with a newborn baby is able to join GenV over a two-year period—potentially up to 150,000 children and their parents."

"Over time, GenV data will be available to approved researchers around the world, so the resulting research could have global significance."

More information: Jian-Rong He et al, Common maternal infections during pregnancy and childhood leukaemia in the offspring: findings from six international birth cohorts, *International Journal of Epidemiology* (2021). [DOI: 10.1093/ije/dyab199](https://doi.org/10.1093/ije/dyab199)

Provided by Murdoch Children's Research Institute

Citation: Common infections in pregnant mums linked to leukemia risk in children (2021, October 28) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-10-common-infections-pregnant-mums-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.