

Development of precision focused ultrasound surgery technology that destructs only the desired tissues

October 28 2021



Illustration of the proposed pressure-modulated shockwave histotripsy pulsing protocol to control the extent and lifetime of a boiling bubble without inducing the shock scattering effect.P1,+ and P1,- are the peak positive and negative pressures in shockwaves at the HIFU focus. P2,+ and P2,- are those in the subsequent HIFU waves within a single pressure-modulated shockwave



histotripsy pulse. Credit: Korea Institute of Science and Technology (KIST)

There are ongoing efforts to use ultrasonic effects with non-invasive properties to replace existing surgical procedures. Focused ultrasound-based bio-tissue destruction technology, which can mechanically fractionate the surrounding bio-tissue using a powerful bubble (cavitation) generated at a short time of approximately 1/100th of a second at the ultrasound focal point, has the advantage of allowing real-time cavitation analysis to monitor the treatment process and a shorter treatment time than the traditional HIFU method, which burns tissues with heat.

According to the Korea Institute of Science and Technology, Dr. Pahk, Ki Joo of the Bionics Research Center has developed a new ultrasound surgery technology that can more precisely and finely fractionate biological <u>tissue</u> using pressure-modulated focused ultrasound (KIST, director Yoon Seok Jin). This new technique is known as Pressuremodulated shockwave histotripsy.

Dr. Pahk was the first in academia to discover the principle of <u>shock</u> scattering by a boiling bubble in the existing focused ultrasound tissue destruction technology in which secondary bubbles are simultaneously generated not only at the ultrasound <u>focal point</u> area but also around it . Though existing focused ultrasound surgery technology has the advantage of being able to mechanically destruct biological tissues, the shock scattering effect reduces precision and makes it difficult to remove tissues or tumors that are close to major organs and blood vessels.

Dr. Pahk devised a method to change the acoustic pressure at the ultrasonic focal point to overcome these limitations. He discovered that



changing the focal acoustic pressure momentarily could control the motility of the bubbles without the shock scattering effect, and thus more precisely destroy the biological tissue, immediately after bubbles are generated at the ultrasonic focal point. He performed acoustic simulations, high speed camera experiments with tissue phantoms, and <u>animal experiments</u> to demonstrate its feasibility.

It was confirmed that by adjusting the acoustic pressure, vapor bubbles with sizes ranging from tens to hundreds of microns could be generated at the ultrasonic focal point without the effect of shock scattering, and that the bubbles could be controlled and adjusted to last for a set period of time. According to the simulation performed by the research team, the reason for this is that the scattered acoustic pressure is lower than the pressure at which the shock scattering effect begins to occur. During the animal testing stage, it was also discovered that bio-tissue can be finely fractionated with a high degree of precision, which is far more precise than existing focused ultrasound technology at the focal point.

Dr. Pahk of KIST explained, "The ultrasound technology we have currently developed is a new technology that can precisely destroy tissues by controlling the size and duration of vapor bubbles. In addition to compensating for the low precision caused by the shock scattering effect, which is the biggest drawback of existing focused ultrasound technology, through bubble motion and duration control, it is possible to adjust the fractionation range and strength. We expect that it will be possible to select and disrupt only specific cells of interest, or that the application will be expanded to decellularization-based cell transplantation research. The relevant core ultrasound <u>technology</u> has been patented in Korea and the United States, and further research is underway to commercialize handheld type <u>ultrasound</u> medical devices capable of precision surgery and treatment."

More information: Ki Joo Pahk, Control of the dynamics of a boiling



vapour bubble using pressure-modulated high intensity focused ultrasound without the shock scattering effect: A first proof-of-concept study, *Ultrasonics Sonochemistry* (2021). DOI: <u>10.1016/j.ultsonch.2021.105699</u>

Provided by National Research Council of Science & Technology

Citation: Development of precision focused ultrasound surgery technology that destructs only the desired tissues (2021, October 28) retrieved 26 April 2024 from https://medicalxpress.com/news/2021-10-precision-focused-ultrasound-surgery-technology.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.