

Prevalence of binge drinking in past 30 days 16.6 percent in 2018

October 18 2021



(HealthDay)—The prevalence of binge drinking within the previous 30



days was 16.6 percent in 2018, with the highest prevalence among those aged 25 to 34 years, according to research published in the Oct. 15 issue of the U.S. Centers for Disease Control and Prevention *Morbidity and Mortality Weekly Report*.

Michele K. Bohm, M.P.H., from the CDC in Atlanta, and colleagues estimated past 30-day binge <u>drinking</u> prevalence, frequency, and intensity overall and by select characteristics and state using data from the 2018 Behavioral Risk Factor Surveillance System.

The researchers found that the overall unadjusted prevalence of binge drinking within the past 30 days was 16.6 percent, representing an estimated 38.5 million U.S. adults. The highest prevalence (26.0 percent) was seen among those aged 25 to 34 years. Compared with women, men had a higher age-standardized prevalence (22.5 versus 12.6 percent), which also increased with income and was highest among non-Hispanic White adults and adults in the Midwest census region. The prevalence of state-level age-standardized binge drinking varied from 10.5 percent in Utah to 25.8 percent in Wisconsin. Twenty-five percent of adults who reported binge drinking did so at least weekly, on average; 25.0 percent consumed at least eight drinks on an occasion.

"Fewer than one half of adults who report binge drinking to a <u>health care</u> <u>provider</u> during a medical checkup are advised to reduce their drinking," the authors write. "Clinicians should follow the U.S. Preventive Services Task Force recommendation to screen all adults for alcohol misuse and provide brief intervention and referral to treatment as needed."

More information: <u>Abstract/Full Text</u>

Copyright © 2021 <u>HealthDay</u>. All rights reserved.



Citation: Prevalence of binge drinking in past 30 days 16.6 percent in 2018 (2021, October 18) retrieved 25 April 2024 from https://medicalxpress.com/news/2021-10-prevalence-binge-days-percent.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.