


# Protein lactylation is induced in the brain by neural activation and social stress

October 12 2021



This paper shows that protein lactylation in brain cells is regulated by neural activation and social defeat stress, paralleled with changes in lactate levels. The stress specifically enhances lysine lactylation of histone H1 proteins in neurons in the prefrontal cortex (PFC). Credit: Hagihara H, et al. Cell Rep 2021

Being bullied leaves a biochemical footprint in the brain. Protein lactylation is a very recently discovered post-translational modification of protein that involves the addition of a lactyl group to lysine residues. Researchers at the Institute for Comprehensive Medical Science at Fujita Health University in Japan, along with colleagues from Ibaraki University and the Japanese National Institute of Advanced Industrial Science and Technology, have now revealed in mice that protein lactylation occurs in neurons in the brain and is positively correlated with lactate levels. This modification was enhanced by neural excitation and social stress, processes known to increase lactate. The stress, from exposure to aggression, specifically enhanced lysine lactylation of histone H1 proteins. The study was published on October 12th in the journal *Cell Reports*.

Lactate (lactic acid) is an end-product of the glycolytic pathway, the rate of which increases along with energy demand. Bodily [lactate](#) concentrations, including in the [brain](#), are enhanced after exercise but also in relation to other conditions. Enhanced levels of brain lactate have been observed in multiple [neuropsychiatric disorders](#), including schizophrenia, bipolar disorder, major depression, and [anxiety disorders](#).

Recent research discovered lysine lactylation as a novel post-translational modification in macrophages that can be stimulated by lactate. The Japanese researchers were interested in whether [protein](#) lactylation also occurs in brain cells.

By biochemistry, cell culture, histology, and mouse behavior experiments, the researchers demonstrated neural activity-dependent protein lactylation in neurons in the mouse brain. *In vivo*, the brain region in which this could be unequivocally established was the prefrontal cortex (PFC). Protein lactylation in the PFC was enhanced after increases in lactate (by intraperitoneal lactate injection), electrical stimulation (by electroconvulsive treatment), anxiety (by exposure to a new open environment), and social stress (by exposure to aggression). The PFC is a brain region involved in top-down control of behavior and emotions.

The [social stress](#) experiment consisted of positioning a mouse for 10 minutes in a cage of a bigger, aggressive mouse, to be vigorously dominated. This was repeated for 10 consecutive days, positioning the poor mouse each day in the cage of another big, aggressive mouse, after which the measurements were made.

"Interestingly, this kind of stress increased brain lactate levels as well as anxiety-like behaviors even after the end of the exposure to the stress, which may cause a decrease of brain pH, a common characteristic among animal models of multiple neuropsychiatric diseases," explains Professor Tsuyoshi Miyakawa, the corresponding author of the study. "Lactate is believed to have an antidepressant effect, and the chronically increased brain lactate levels may form a compensatory mechanism against stress."

By employing [mass spectrometry](#), the researchers identified 63 lactylated proteins in the mouse PFC. Among these proteins, lactylation of H1 histones was significantly upregulated after exposure of mice to aggression. Super-resolution imaging with stimulated emission depletion (STED) microscopy revealed that the stress increased lactylation of histone H1 in the nuclei of neurons, but not astrocytes.

Histone modifications mediate epigenetic regulation of a wide variety of genes. Dr. Hideo Hagihara, lead author of the paper, says: "The lactate-sensitive epigenetic changes in the context of stress may play a role in gene-mediated regulation of behavior. However, for now, these data remain entirely correlational. It is important to provide mechanistic insights on how histone lactylation may affect [stress](#)-related responses. We are currently conducting experiments to address the issue."

**More information:** Hagihara H. et al. Protein lactylation induced by neural excitation. *Cell Reports* (2021). doi: [doi.org/10.1016/j.celrep.2021.109820](https://doi.org/10.1016/j.celrep.2021.109820)

Hideo Hagihara et al, Systematic analysis of brain lactate and pH levels in 65 animal models related to neuropsychiatric conditions, *bioRxiv* (2021). [DOI: 10.1101/2021.02.02.428362](https://doi.org/10.1101/2021.02.02.428362)

Provided by Fujita Health University

Citation: Protein lactylation is induced in the brain by neural activation and social stress (2021, October 12) retrieved 5 May 2024 from <https://medicalxpress.com/news/2021-10-protein-lactylation-brain-neural-social.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--