


Researchers reveal mechanism of oligodendrocyte myelination by osteocalcin hormone

October 22 2021



A schematic depicting the action of OCN on myelin regulation in oligodendrocytes. Credit: SIAT

Osteocalcin (OCN) is a multifunctional bone-derived hormone that modulates numerous physiological activities. OCN can cross the blood brain barrier (BBB) and thus play critical roles in neuronal and brain development.

Fully elucidating the exact function of OCN in the [central nervous system](#) (CNS) is significant for the discovery of therapeutic targets in treating CNS disorders.

A research team led by Li Xiang from the Shenzhen Institute of Advanced Technology (SIAT) of the Chinese Academy of Sciences has revealed the molecular mechanism of oligodendrocyte [myelination](#) by OCN in the CNS.

Their study was published in *Science Advances* on Oct. 22.

The researchers first investigated the effects of OCN on CNS myelination using an OCN-deficient mouse model. By combining immunostaining, western blotting and electron microscopy analysis, they found that OCN deletion results in hypermyelination in the CNS.

Oligodendrocytes (OLs) are [glial cells](#) that form myelin in the CNS. To clarify the mechanism of OCN in CNS myelination, the researchers further investigated whether OCN plays a role in regulation of OL differentiation and myelination.

By using an OCN-deficient mouse model, primary OL culture and exogenous OCN treatment, they found that genetic deletion of OCN facilitated OL differentiation and hypermyelination through the transcription factor *Myrf* in the CNS. Although dispensable for the proliferation of oligodendrocyte precursor cells (OPCs), OCN was critical for the myelination of OLs, which affected myelin production and remyelination after demyelinating injury.

To identify the receptor for OCN in OLs, the researchers combined RNA sequencing, heterologous systems biophysical analysis, and *in vivo* mouse model validation, and finally proved that GPR37 is the receptor for OCN and mediates its function in OLs.

These findings not only identify a new role for the OCN hormone in the CNS, but also demonstrate a [molecular basis](#) for the function of OCN by activation of GPR37 signaling. This study provides a [unique perspective](#)

for understanding the mutual communication between skeletal biology and glial functions.

More information: Zhengjiang Qian et al, Osteocalcin attenuates oligodendrocyte differentiation and myelination via GPR37 signaling in the mouse brain, *Science Advances* (2021). [DOI: 10.1126/sciadv.abi5811](https://doi.org/10.1126/sciadv.abi5811) . www.science.org/doi/10.1126/sciadv.abi5811

Provided by Chinese Academy of Sciences

Citation: Researchers reveal mechanism of oligodendrocyte myelination by osteocalcin hormone (2021, October 22) retrieved 26 June 2024 from <https://medicalxpress.com/news/2021-10-reveal-mechanism-oligodendrocyte-myelination-osteocalcin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.