

Study first to reveal specific molecular mechanism that controls the transition from acute to chronic pain

October 22 2021



"This study is the first to identify that NAAA, a previously unrecognized control node, can be effectively targeted by small-molecule therapeutics that inhibit this enzyme, and block the transition from acute to chronic pain," said Daniele Piomelli, Ph.D., Distinguished Professor in the UCI School of Medicine Department of Anatomy & Neurobiology. Credit: UCI School of Medicine

A new study led by University of California, Irvine researchers is the first to reveal the specific molecular mechanism that controls the transition from acute to chronic pain, and identifies this mechanism as a critical target for disease-modifying medicines.

Findings from the study, titled "NAAA-regulated lipid signaling governs the transition from acute to [chronic pain](#)," published today in *Science Advances*, show that disabling N-acylethanolamine acid amidase (NAAA)—an intracellular enzyme—in the [spinal cord](#) during a 72-hour time window following peripheral tissue injury halts chronic pain development in male and female mice.

"Delineating the nature, localization and timing of the events involved in pain chronicity is necessary to pinpointing control nodes in the process that can be targeted by new classes of disease-modifying medicines beyond analgesics," said Daniele Piomelli, Distinguished Professor in the UCI School of Medicine Department of Anatomy & Neurobiology.

"This study is the first to identify that NAAA, a previously unrecognized control node, can be effectively targeted by small-molecule therapeutics that inhibit this enzyme, and block the transition from acute to chronic pain."

Chronic pain evolves from [acute pain](#) caused by the physical trauma of tissue damage due to surgery or injury and is a massive problem, affecting more than 1.5 billion people worldwide. Chronic pain continues long past tissue healing, is often resistant to therapy, and remains seriously undertreated. Treatment is largely dependent on a handful of analgesic drug classes such as opioids, which may lose effectiveness over time and can also lead to addiction. Nerve damage is considered to be a critical factor in the transition to chronic pain, but the underlying molecular events leading to its emergence have been poorly understood.

"Our findings suggest a new class of drugs—NAAA inhibitors—can be used to treat various forms of chronic [pain](#) and in preventing incisional and inflammatory injuries following surgery," Piomelli said.

More information: Yannick Fotio et al, NAAA-regulated lipid signaling governs the transition from acute to chronic pain, *Science Advances* (2021). DOI: [10.1126/sciadv.abi8834](https://doi.org/10.1126/sciadv.abi8834).
www.science.org/doi/10.1126/sciadv.abi8834

Provided by University of California, Irvine

Citation: Study first to reveal specific molecular mechanism that controls the transition from acute to chronic pain (2021, October 22) retrieved 20 March 2024 from <https://medicalxpress.com/news/2021-10-reveal-specific-molecular-mechanism-transition.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--