


Identifying specific hazard patterns for emerging cardiovascular disease

October 18 2021



Credit: © U. Flögel

Predicting when atherosclerotic changes in the arteries in a stable state will progress to acute cardiovascular disease has remained unresolved.

The authors of the paper currently published in *Nature Communications*, led by first author Prof. Ulrich Flögel, MD, Institute of Molecular Cardiology, Faculty of Medicine, Heinrich Heine University, Düsseldorf, Germany, present an imaging technique—the targeted and multicolor nanotracer platform technology—that visualizes the hazard patterns in the development of progressive coronary disease in a mouse model.

Hazard pattern using the example of a mouse heart (top in cross-section) with a massively worn out and thus in its performance strongly impaired right heart as consequential damage to be seen in the follow-up examination (bottom right compared to bottom left).

The cascade of these vascular diseases ranges from inflammation of the vessel, thrombosis with subsequent detachment of minute particles of the vascular plaque, to vascular occlusion resulting in permanent damage due to, for example, myocardial infarction.

To show the hazard patterns, three molecules (ligands)—are coupled to different types of perfluorocarbon nanoemulsion. They are specifically directed respectively to sites of inflammation, acute and chronic thrombi. There, they embed themselves and become visible and distinguishable through the use of a specific MRI [imaging technique](#) (¹⁹F-MRI).

This enables the scientists to identify areas that are at high risk of developing a heart attack. This happens at a time when no conventional parameter in laboratory diagnostics or ECG indicates an imminent danger. The research team shows that these patterns can be visualized reliably—and before the onset of infarction—by the "multi-color multi-targeted ¹H/¹⁹F MRI" technique they developed. Only a single MRI scan is required, which can localize individually affected sites and differentiate them by color with respect to inflammation, freshly

developed or advanced thrombi.

Thereby, the early identified vulnerable sites may well be different in the individual ([mouse model](#)). However, it is important that in each case the corresponding massive secondary damage also occurred at these sites. In further studies, Flögel et al. showed that the "multi-targeted 1H/19F MRI method" is able to clearly visualize the entire vicious circle of platelet adhesion, infiltration of immune cells and formation of thrombi leading to coronary atherothrombosis, [myocardial infarction](#) and severe deterioration of ventricular function.

More information: Ulrich Flögel et al, Multi-targeted 1H/19F MRI unmasks specific danger patterns for emerging cardiovascular disorders, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-26146-6](https://doi.org/10.1038/s41467-021-26146-6)

Provided by Heinrich-Heine University Duesseldorf

Citation: Identifying specific hazard patterns for emerging cardiovascular disease (2021, October 18) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-10-specific-hazard-patterns-emerging-cardiovascular.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--