


# How can the brain perceive familiar objects when they become indistinct

November 26 2021



We can perceive familiar objects even when they become indistinct (A). After repeatedly encountering the objects, a subset of neurons in the primary visual cortex exhibits strong responses to low-contrast visual stimuli. These low-contrast-preferring neurons substantially contribute to low-contrast visual information representation, ensuring a consistent perception of familiar objects with contrast variation (B). Credit: Rie Kimura

Researchers from NIPS explored the brain neuronal mechanism that allows the perception of familiar images even if they are indistinct. They found that the number of neurons responding to low-contrast rather than high-contrast visual stimuli increased in rats performing a visual orientation discrimination task after repeated experiences. These neurons showed stronger activities in correct-choice than incorrect-choice trials. These neurons efficiently represented low-contrast stimulations. Thus, the low-contrast preference in V1 activity may contribute to improved low-contrast visual discrimination.

The appearance of objects can often change. For example, in dim evenings or fog, the contrast of the objects decreases, making it difficult to distinguish them. However, after repeatedly encountering specific objects, the brain can identify them even if they become indistinct. The exact mechanism contributing to the perception of low-contrast familiar objects remains unknown.

In the [primary visual cortex](#) (V1), the area of the cerebral cortex dedicated to processing basic visual information, the visual responses have been considered to reflect directly the strength of external inputs. Thus, high-contrast visual [stimuli](#) elicit strong responses and vice versa.

In this study, Rie Kimura and Yumiko Yoshimura found that in rats, the number of V1 neurons preferentially responding to low-contrast stimuli

increases after repeated experiences. In these neurons, low-contrast visual stimuli elicit stronger responses, and high-contrast stimuli elicit weaker responses. These low contrast–preferring neurons show a more evident activity when rats correctly perceive a low-contrast familiar [object](#). It was first reported in *Science Advances* that low-contrast preference in V1 is strengthened in an experience-dependent manner to represent low-contrast visual information well. This mechanism may contribute to the perception of familiar objects, even when they are indistinct.

"This flexible information representation may enable a consistent perception of familiar objects with any contrast," Kimura says. "The flexibility of our brain makes our sensation effective, although you may not be aware of it. An artificial neural network model may reproduce the human sensation by incorporating not only [high contrast](#)–preferring neurons, generally considered until now, but also low contrast–preferring [neurons](#), the main focus of this research."

**More information:** Rie Kimura, The contribution of low contrast-preferring neurons to information representation in the primary visual cortex after learning, *Science Advances* (2021). [DOI: 10.1126/sciadv.abj9976](#). [www.science.org/doi/10.1126/sciadv.abj9976](http://www.science.org/doi/10.1126/sciadv.abj9976)

Provided by National Institute for Physiological Sciences

Citation: How can the brain perceive familiar objects when they become indistinct (2021, November 26) retrieved 2 May 2024 from <https://medicalxpress.com/news/2021-11-brain-familiar-indistinct.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--