


# Brain stimulation could help you decide when to give up

November 18 2021

---



Credit: AI-generated image ([disclaimer](#))

New methods of brain stimulation are being tested to see whether they can be used to enhance cognitive behavior. In this study, researchers from Japan have found that stimulation at specific frequencies can modulate thought processes related to problem solving.

In a study published this month in *Scientific Reports*, researchers from the University of Tsukuba have revealed that a non-invasive form of [brain](#) stimulation called [repetitive transcranial magnetic stimulation](#), or rTMS, can be used to modulate [brain rhythms](#) and [cognitive behavior](#) related to "giving-up" during [problem solving](#).

rTMS has been found to increase the synchronization of brain rhythms during cognitive tasks. Because specific brain rhythms are related to aspects of cognition, this could have great potential in helping people to enhance their cognitive abilities, something the researchers at the University of Tsukuba aimed to address.

"Prefrontal theta rhythms are known to be associated with cognitive control and conflict resolution, while alpha rhythms are related to the generation of new ideas during problem solving," says lead author of the study Professor Masahiro Kawasaki. "Because rTMS can be used to alter [neural activity](#), we wanted to investigate how stimulation at specific frequencies could affect problem-solving behavior."

To do this, the researchers examined the relationship between brain oscillations and cognitive "giving-up" while participants performed problem-solving tasks. Then, they examined the effects of rTMS on brain oscillations and task performance. The task involved solving riddles, and the participants were allowed to indicate when they "gave up" on the task. "We found the theta rhythm in the frontal lobe of the brain was associated with "giving-up," while the alpha [rhythm](#) was associated with successful problem solving," explains Professor Kawasaki.

Next, the researchers applied rTMS at specific frequencies while the participants completed the same cognitive task, and measured brain waves and task performance.

"The results were exciting," says Professor Kawasaki. "Theta-frequency rTMS increased theta amplitudes and decreased "giving-up" behaviors, and alpha-frequency rTMS increased alpha amplitudes but had no effect on "giving-up." This is important evidence that rTMS can be used to modulate brain oscillatory activity and behaviors related to "giving-up" processes."

During rumination, which is a common feature of depression, adaptive "giving-up" behaviors can be suppressed. For instance, individuals may have trouble deciding to "give up" and focus on something else when a certain behavior is not leading to the desired outcome. This research shows how modulating neural rhythms such as alpha and theta could decrease rumination, and thus treat symptoms of depression. Furthermore, this [study](#) is an important example of how rTMS can be used to change behavior related to cognitive performance via the modulation of task-related brain rhythms.

**More information:** Eri Miyauchi et al, Behavioural effects of task-relevant neuromodulation by rTMS on giving-up, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-01645-0](https://doi.org/10.1038/s41598-021-01645-0)

Provided by University of Tsukuba

Citation: Brain stimulation could help you decide when to give up (2021, November 18) retrieved 23 June 2024 from <https://medicalxpress.com/news/2021-11-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--