


# Diabetes drug slows kidney function decline

November 6 2021

---



Credit: Pixabay/CC0 Public Domain

Clinical trial data reveal that dapagliflozin— a sodium-glucose cotransporter 2 inhibitor prescribed to treat diabetes—reduces the rate of kidney function decline in patients with chronic kidney disease (CKD). The findings will be presented at ASN Kidney Week 2021 November 4–November 7.

The DAPA-CKD trial randomized 4,304 participants with CKD to dapagliflozin 10 mg or placebo once daily, added to standard care.

Although participants without diabetes also experienced a slower rate of kidney function decline with dapagliflozin, the effect of dapagliflozin was greater in those with diabetes.

"The key conclusion is that dapagliflozin is an [effective treatment](#) to slow progressive kidney function loss in patients with CKD with and without type 2 diabetes," said lead author Hiddo Lambers Heerspink, Ph.D., PharmD, of the University Medical Center Groningen.

"Therefore, in addition to reducing the risk of heart failure or mortality, as previously shown in the DAPA-CKD trial, dapagliflozin also slows the progression of kidney function decline."

Study: "The effect of dapagliflozin on rate of kidney function decline in patients with [chronic kidney disease](#): a prespecified analysis from the DAPA-CKD"

Provided by American Society of Nephrology

Citation: Diabetes drug slows kidney function decline (2021, November 6) retrieved 27 April 2024 from <https://medicalxpress.com/news/2021-11-diabetes-drug-kidney-function-decline.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.